

THE UNIVERSITY OF CHICAGO

DETERMINING WHETHER CERTAIN AFFINE DELIGNE-LUSZTIG SETS
ARE EMPTY

A DISSERTATION SUBMITTED TO
THE FACULTY OF THE DIVISION OF THE PHYSICAL SCIENCES
IN CANDIDACY FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

DEPARTMENT OF MATHEMATICS

BY
DANIEL C. REUMAN

CHICAGO, ILLINOIS
AUGUST 2002

ABSTRACT

Let F be a non-archimedean local field, let L be the maximal unramified extension of F , and let σ be the Frobenius automorphism. Let G be a split connected reductive group over F , and let $\mathcal{B}_\infty$ be the Bruhat-Tits building associated to $G(L)$. Let $\mathcal{B}_1$ be the building associated to $G(F)$. Then σ acts on $G(L)$ with fixed points $G(F)$. Let I be the Iwahori associated to a chamber in $\mathcal{B}_1$. We have the relative position map $inv : G(L)/I \times G(L)/I \rightarrow \tilde{W}$, where $\tilde{W}$ is the extended affine Weyl group of G . If $w \in \tilde{W}$ and $b \in G(L)$, then the affine Deligne-Lusztig set $X_w(b\sigma)$ is $\{x \in G(L)/I : inv(x, b\sigma(x)) = w\}$. This dissertation answers the question of which $X_w(b\sigma)$ are non-empty for certain G and b .

ACKNOWLEDGEMENTS

Above all, I would like to thank my advisor, Robert Kottwitz, for being extremely helpful at every turn. Professor Kottwitz's mathematical guidance has helped with every aspect of this dissertation. His support and understanding regarding non-research issues encountered during graduate education is also remarkable. I am very grateful to have been his student. I would also like to thank Chris Degni and Sudheer Shukla for helpful discussions and lectures. Finally, I thank Moon Duchin, Kaj Gartz, Mark Behrens, and Moses Hohman for moral support.

TABLE OF CONTENTS

ABSTRACT	ii
ACKNOWLEDGEMENTS	iii
LIST OF FIGURES	vi
LIST OF TABLES	x
1 INTRODUCTION	1
2 GROUPS OF SEMISIMPLE RANK 1	5
2.1 SL_2	5
2.1.1 Identity σ -conjugacy Class	5
2.1.2 Non-identity σ -conjugacy Classes	7
2.1.3 Symmetry Under a $\mathbb{Z}/2$ -action	8
2.1.4 Relationship Between $X_w(1\sigma)$ and $X_w(b\sigma)$	10
2.2 GL_2 and PGL_2	13
2.3 Reorganization of Results	15
3 GROUPS OF SEMISIMPLE RANK 2	18
3.1 SL_3	18
3.1.1 Standard Minimal Galleries and Composite Galleries	19
3.1.2 A Superset of the Solution Set	21
3.1.3 Relationship Between $X_w(1\sigma)$ and $X_w(b\sigma)$, and an Efficient Way of Computing Supersets	50
3.1.4 A Method Suggested by Rapoport and Kottwitz	55
3.1.5 A Subset of the Solution Set for $b = 1$	59
3.1.6 A Geometric Construction of a Subset of the Solution Set	66
3.1.7 Symmetry Under a $\mathbb{Z}/3$ -action	95
3.2 GL_3 and PGL_3	96
3.3 Sp_4	97
3.3.1 Standard Minimal Galleries and Composite Galleries	98
3.3.2 A Conjectural Superset of the Solution Set	100
3.3.3 Relationship Between $X_w(1\sigma)$ and $X_w(b\sigma)$ for Sp_4 , and an Ef- ficient Way of Computing Supersets	106
3.3.4 The Method of Kottwitz and Rapoport Applied to Sp_4	106
3.3.5 A Subset of the Solution Set for $b = 1$ for Sp_4	110

3.3.6	Subsets for Other b , and Symmetry under a $\mathbb{Z}/2$ -action	112
3.4	Comments on GSp_4 , PSp_4 , and G_2	119
3.5	Invariance Properties of Solution Sets	120
REFERENCES		125

LIST OF FIGURES

2.1	Chamber labels	6
2.2	Alternate chamber labels	6
2.3	Proof of Proposition 2.1.2	7
2.4	Summary of SL_2 results	9
2.5	Lattice labels for vertices	10
2.6	For w in the positive Weyl chamber	11
2.7	For w in the negative Weyl chamber	12
2.8	Summary of some results for GL_2	14
2.9	Computations for some σ -conjugacy classes for GL_2	15
2.10	Summary of some more results for GL_2	15
3.1	Primary and secondary directions	20
3.2	Example SMG type and edge of departure	22
3.3	Composite gallery example	22
3.4	Results from the SMG in Figure 3.2	23
3.5	Γ_E^1 possibilities for I_1	24
3.6	Some composite gallery possibilities for I_1	25
3.7	Class I_1 , where Γ_E^1 has 3 chambers after the turning edge	26
3.8	Class I_1 , where Γ_E^1 has 5 chambers after the turning edge	27
3.9	Class I_1 , where Γ_E^1 has 7 chambers after the turning edge	28
3.10	Summary of results for I_1 , an odd number of chambers after the turning edge	29
3.11	More composite gallery possibilities for I_1	30
3.12	Class I_1 , where Γ_{E_1} has 2 chambers after the turning edge	31
3.13	Class I_1 , where Γ_{E_1} has 4 chambers after the turning edge	32
3.14	Class I_1 , where Γ_{E_1} has 6 chambers after the turning edge	33
3.15	Summary of results for I_1 , an even number of chambers after the turning edge	34
3.16	Some composite gallery possibilities for I_2	35
3.17	More composite gallery possibilities for I_2	35
3.18	Class I_2 , where Γ_{E_1} has 0, 1 or 2 chambers after the departure edge, $w = 1$	35
3.19	Class I_2 , where Γ_{E_1} has 3 or 4 chambers after the departure edge, $w = 1$	36
3.20	Class I_2 , where Γ_{E_1} has 5 or 6 chambers after the departure edge, $w = 1$	36
3.21	Summary of results for I_2 , $w = 1$	37
3.22	Summary of results for I_2 , $w = f$	38
3.23	Summary of results for I_1 and I_2	39

3.24	Result for $\alpha = 2, \beta = 0$	41
3.25	Result for $\alpha = 2, \beta = 1$	42
3.26	Result for $\alpha = 4, \beta = 0$	43
3.27	Result for $b = 1$	44
3.28	General shape of composite galleries for $b = 1, w = f, I_1$	45
3.29	General shape of composite galleries for $b = 1, w = 1, I_1$	45
3.30	General shapes of composite galleries for $b = 1, w = 1, I_2$	46
3.31	Result for $\alpha = 4, \beta = -2$	47
3.32	bC_M for degenerate b	48
3.33	General shape of composite galleries for $b \neq 1$ with $\alpha + 2\beta = 0, w = f, I_1$	49
3.34	General shape of composite galleries for $b \neq 1$ with $\alpha + 2\beta = 0, w = 1, I_1$	49
3.35	$\Omega_1^2, \Omega_f^2, \Omega_{p,1}^1$ and $\Omega_{p,f}^1$	53
3.36	Choice points for a sample composite gallery	54
3.37	Choice points for $\Omega_{p,1}^1$ and Γ_x	54
3.38	How to address choice points of Γ_x that occur before the turning point of Γ_E^3	55
3.39	Main results of the method suggested by Rapoport and Kottwitz	57
3.40	The $a = 1$ results of the method suggested by Rapoport and Kottwitz	58
3.41	Results from the method suggested by Rapoport and Kottwitz under the circumstances of Lemma 3.1.5	58
3.42	The choice D_1 of xC_M	59
3.43	The resulting Γ_x for D_1 and a particular choice of aw	60
3.44	The result of folding galleries of the same type as $x^{-1}\Gamma_x$ for the Γ_x pictured in Figure 3.43	60
3.45	Results for $xC_M = D_1$ and $w = r^2$ or for $xC_M = D_1$ and $w = r$	61
3.46	The choices D_2 and D_3 of xC_M	62
3.47	Results for $xC_M = D_2$ and $w = r^2$ or for $xC_M = D_2$ and $w = r$	63
3.48	Results for $xC_M = D_3$ and $w = r^2$ or for $xC_M = D_3$ and $w = r$	64
3.49	Combination of all results referred to in Table 3.1	65
3.50	Some choices for xC_M that are two chambers away from C_M	65
3.51	A Useful Lemma	66
3.52	An example of P, P_b and P_σ	68
3.53	w_2, w_3, w_4 and w_5	68
3.54	Multiple rows of the w_i	69
3.55	Filling in the rest of $\overline{P}$	69
3.56	The definition of the invariant γ_2 , part 1	70
3.57	The definition of the invariant γ_2 , part 2	71
3.58	The definition of the invariant γ_2 , part 3	71
3.59	$T(P) = S(P) = \infty$	72
3.60	The meaning of a tilde	73

3.61	Existence of all values of γ_1 and γ_2 , case 1	74
3.62	Existence of all values of γ_1 and γ_2 , case 2 subcase 1	76
3.63	Existence of all values of γ_1 and γ_2 , case 2 subcase 2	77
3.64	Existence of all values of γ_1 and γ_2 , case 2 subcase 3	78
3.65	Existence of all values of γ_1 and γ_2 , case 2 subcase 4	80
3.66	Definition of negative values of γ_2	81
3.67	Existence of all negative values of γ_2 , case 1	82
3.68	Existence of all negative values of γ_2 , case 2 subcase 1	84
3.69	Existence of all negative values of γ_2 , case 2 subcase 1 diagram 2 . . .	85
3.70	Existence of all negative values of γ_2 , case 2 subcase 2	86
3.71	$x^{-1}(\overline{P} \cup P_\sigma)$ for $\gamma_2 > 0$	87
3.72	Example of how γ_1 and γ_2 determine $\rho(x^{-1}b\sigma(x)C_M)$	88
3.73	$x^{-1}(\overline{P} \cup P_\sigma)$ for $\gamma_2 \leq 0$	89
3.74	$x^{-1}(\overline{P} \cup P_\sigma)$ for $\gamma_2 \leq 0$, folding explained	90
3.75	$x^{-1}(\overline{P} \cup P_\sigma)$ for $\gamma_2 \leq 0$, folding explained further	91
3.76	Choice Points	92
3.77	A composite gallery for $p = 1$, $\alpha = 2$, $\beta = -1$	93
3.78	All foldings of composite galleries with $p = 1$ occur, where $b \neq 1$ and $\alpha + 2\beta = 0$	94
3.79	Definition of γ_3	95
3.80	What happens to Figure 3.79 when $\gamma_3 = 0$	96
3.81	How the torus of Sp_4 acts on A_M	98
3.82	Primary and Secondary Directions in Sp_4	99
3.83	Result for $b = 1$, Sp_4	101
3.84	Result for $\alpha = 3$, $\beta = 1$, Sp_4	102
3.85	Result for $\alpha = 6$, $\beta = 3$, Sp_4	103
3.86	Result for $\alpha = 6$, $\beta = 5$, Sp_4	104
3.87	Result for $\alpha = 7$, $\beta = 1$, Sp_4	105
3.88	Half infinite galleries for Sp_4 , $\alpha > \beta$ (specifically, $\alpha = 3$, $\beta = 1$)	107
3.89	Half infinite galleries for Sp_4 , $\alpha = \beta$ (specifically, $\alpha = \beta = 3$)	108
3.90	Main results of the method suggested by Rapoport and Kottwitz, Sp_4	109
3.91	The $a = 1$ results of the method suggested by Rapoport and Kottwitz, Sp_4	110
3.92	Results of the method suggested by Rapoport and Kottwitz under the circumstances of Lemma 3.3.4, Sp_4	111
3.93	The choices D_1 , D_2 and D_3 of G_1	112
3.94	Results for $xC_M = D_1$ and $w = r$ or for $xC_M = D_1$ and $w = r^3$. . .	113
3.95	Results for $xC_M = D_2$ and $w = r$ or for $xC_M = D_2$ and $w = r^3$. . .	114
3.96	Results for $xC_M = D_3$ and $w = r$ or for $xC_M = D_3$ and $w = r^3$. . .	115
3.97	Results for $xC_M = D_1$ and $w = r^2$	116
3.98	Results for $xC_M = D_2$ and $w = r^2$	117
3.99	Results for $xC_M = D_3$ and $w = r^2$	118

3.100	The translations ϵ and δ of A_M for G_2	120
3.101	Region R	123
3.102	Understood region after applying invariance properties to region R .	124

LIST OF TABLES

2.1	$D_M^{\pm i}$ intersects exactly these Σ_j non-trivially for i even	16
2.2	$D_M^{\pm i}$ intersects exactly these Σ_j non-trivially for i odd	16
2.3	σ -conjugacy classes that intersect (D_M^i, j) non-trivially	17
3.1	Which figure contains which results	62

CHAPTER 1

INTRODUCTION

Let F be a non-archimedean local field with ring of integers $\mathcal{O}_F$, and let G be a split connected reductive group over F . Let L be the completion of the maximal unramified extension of F . Let σ be the Frobenius automorphism of L over F . Let $\mathcal{B}_n$ be the affine building for $G(E)$ where E/F is the unramified extension of degree n in L , and let $\mathcal{B}_\infty$ be the affine building for $G(L)$. The theory of buildings was developed by Bruhat and Tits in [1] and [2], and is reviewed by Garrett in [4]. We know that σ acts on $\mathcal{B}_\infty$, and the fixed points of σ^n are $\mathcal{B}_n$. Let T be a split torus in G and let I be an Iwahori in $G(L)$ containing $T(\mathcal{O}_L)$, where $\mathcal{O}_L$ is the ring of integers of L . Let A_M and C_M be the correspondingly specified apartment and chamber, which we assume are in $\mathcal{B}_1$. We will call these the *main apartment* and the *main chamber*, respectively.

If $b \in G(L)$ then the σ -conjugacy class of b is $\{x^{-1}b\sigma(x) : x \in G(L)\}$. If $\tilde{W}$ is the extended affine Weyl group, and $w \in \tilde{W}$ then we define, after Rapoport and Kottwitz, the affine Deligne-Lusztig set $X_w(b\sigma) = \{x \in G(L)/I : \text{inv}(x, b\sigma(x)) = w\}$. Here $\text{inv} : G(L)/I \times G(L)/I \rightarrow \tilde{W}$ is the relative position map. Note that if b_1 and b_2 are σ -conjugate then the sets $X_w(b_1\sigma)$ and $X_w(b_2\sigma)$ are in bijective correspondence. When $F = \mathbb{F}_q((t))$, the affine Deligne-Lusztig set can be given variety structure over $\mathbb{F}_q$ (locally of finite type).

Sometimes the $X_w(b\sigma)$ are empty. Our goal is to determine when the $X_w(b\sigma)$ are non-empty. This paper concerns itself mainly with the question of which $X_w(b\sigma)$ are non-empty for groups of semisimple rank 1 and 2.

The question of which $X_w(b\sigma)$ are non-empty for $\{x^{-1}b\sigma(x) : x \in G(L)\}$ a fixed σ -conjugacy class can be rephrased by saying that $X_w(b\sigma)$ is non-empty if and only if $\{x^{-1}b\sigma(x) : x \in G(L)\}$ intersects the double- I -coset corresponding to w in a non-trivial way. Here we are using the standard correspondence between $\tilde{W}$ and $I \backslash G(L) / I$. We will use this correspondence implicitly for the rest of the paper. To see that the above rephrasing is accurate, note first that $\text{inv}(x, b\sigma(x)) = \text{inv}(1, x^{-1}b\sigma(x))$ is the relative position of 1 and $x^{-1}b\sigma(x)$. By definition, $\text{inv}(1, x^{-1}b\sigma(x)) = Ix^{-1}b\sigma(x)I$. We see that $X_w(b\sigma)$ is non-empty if and only if there exists x such that $Ix^{-1}b\sigma(x)I = w$, which is what was to be proven.

If G is simply connected, then $\tilde{W} \simeq W_a$ is in bijective correspondence with chambers in the main apartment A_M . Here W_a is the affine Weyl group of G . So for a fixed σ -conjugacy class $\{x^{-1}b\sigma(x) : x \in G(L)\}$, the set of w for which $X_w(b\sigma)$ is non-empty can be represented geometrically as a collection of chambers in A_M . For

groups of rank 1 and 2, one can therefore draw pictures of the solution set. This is the form in which we present results in this paper.

Complete information can be obtained without difficulty for groups of semisimple rank 1. This is done in Chapter 2. The solution sets for each σ -conjugacy class of SL_2 are pictured in Figure 2.4. The solution sets for each σ -conjugacy class of GL_2 or PGL_2 are pictured in Figures 2.8 and 2.10.

The group of semisimple rank 2 for which the most information has been obtained is $G = SL_3$. We have worked only with σ -conjugacy classes that are basic in the maximal torus. These are listed at the beginning of Section 3.1. The general approach by which we have described $\{inv(x, b\sigma(x)) : x \in SL_3(L)\} \subseteq \tilde{W} \simeq W_a$ in these cases has two main parts. The first part, done in Sections 3.1.1, 3.1.2, and 3.1.3, seeks to produce a superset of the solution set. The general methodology here is to establish a way of choosing, for each $x \in SL_3(L)$, a gallery Γ_x between xC_M and $b\sigma(x)C_M$. Possible values of $inv(x, b\sigma(x))$ can then be enumerated by listing the possible “foldings” of $x^{-1}\Gamma_x$. Optimizations can be implemented so that this is a finite, but still prohibitively lengthy computation. It has only been carried out to completion for two σ -conjugacy classes. The results are pictured in Figure 3.23 and Figure 3.27. In the process of doing these two complete computations, we observe that in fact one need only do a much smaller part of the complete computation to obtain all elements of W_a that arise in the superset. We do not have a proof that this is the case in general, but we are very strongly confident that it is. Section 3.1.3 develops an efficient way of doing the smaller, partial computation that works for any σ -conjugacy class. Results of the smaller computation that are (confidently) conjectured to be the results of the entire computation are listed for various σ -conjugacy classes in Figures 3.24, 3.25, 3.26, and 3.31.

The other half of the work for SL_3 is to develop a subset of the solution set. This is done in two different ways. The first method, which is discussed in Sections 3.1.4 and 3.1.5, is simple and potentially generalizable to higher rank groups, but only applies to the case $b = 1$. The second method (discussed in Section 3.1.6) applies for any b in the collection of σ -conjugacy classes under consideration, but is very complicated and somewhat aesthetically unpleasing. Also, it probably could not be generalized to higher rank groups. In all cases, the subset results turn out to be equal to the superset results discussed above. So it is known that Figures 3.23 and 3.27 are solution sets for their respective σ -conjugacy classes, and we are very confident (although it is not proved) that Figures 3.24, 3.25, 3.26, and 3.31 are solution sets for their respective σ -conjugacy classes. It is known that these last four figures are at least subsets of their respective solution sets.

The same methods can be applied to Sp_4 , although this has been carried out to a smaller extent. Again, we only consider σ -conjugacy classes b which are basic in the maximal torus. The complete computation of the superset of the solution set has only been done for $b = 1$. The result is pictured in Figure 3.83. Again, there is a partial computation which gives all results obtained from the complete computation

in this case. Carrying out the partial computation for some other σ -conjugacy classes (done in Section 3.3.2) yields Figures 3.84, 3.85, 3.86, and 3.87. We conjecture that these are complete supersets, but we are not completely confident of this because the chambers marked with a * seem to be holes in the pattern.

The subset methods of Section 3.1.4 and 3.1.5 have been applied to Sp_4 , $b = 1$ in Sections 3.3.4 and 3.3.5. The result is the same as the superset pictured in Figure 3.83, so this figure gives the solution set for $b = 1$. The complicated methods of Section 3.1.6 have not been adapted to Sp_4 , although such an adaptation probably could be done. We conjecture that one would obtain subsets equal to the sets pictured in Figures 3.84, 3.85, 3.86, and 3.87, making these figures actual solution sets. So for Sp_4 we have definite results for $b = 1$, and conjectural results of which we are not completely confident for other b .

The results obtained for SL_3 can easily be adapted to GL_3 and PGL_3 . This is done in Section 3.2. Similarly, the results for Sp_4 can be adapted to GSp_4 and PSp_4 . This is done in Section 3.4. The methods used could be applied to G_2 as well. This is also discussed in Section 3.4.

We also say a few words describing the basic ideas behind the methods used. The backbone of the method by which supersets of solution sets are produced is the “folding” of galleries. If G is a gallery starting at C_M , then the *folding results* of G are the possible values of the last chamber of $\rho(\tilde{G})$, where $\rho : \mathcal{B}_\infty \rightarrow A_M$ is the retraction centered at C_M , and $\tilde{G}$ is any gallery of the same “shape” as G that starts at C_M . To produce a superset, it turns out that the folding results have to be calculated for a gallery associated to each chamber in $\mathcal{B}_\infty$. Perhaps the real content of the method is how these computations can be grouped so that they can be done efficiently. This will be discussed in Sections 3.1.2 and 3.1.3 for SL_3 , and Sections 3.3.2 and 3.3.3 for Sp_4 .

As mentioned previously, there are two methods used to produce subsets of solution sets. The first, which only applies for $b = 1$, is carried out in Sections 3.1.4 and 3.1.5 for SL_3 , and Sections 3.3.4 and 3.3.5 for Sp_4 . This method gets started using an observation of Kottwitz and Rapoport that provides, with very little work, a large and well distributed collection of chambers which must be contained in the solution set. This collection can then be enlarged as follows. If $inv(x, \sigma(x)) = w$ is one of the chambers provided by Kottwitz’s and Rapoport’s observation, and if Γ is a minimal gallery between xC_M and $\sigma(x)C_M$, then Γ describes the relative position w . One can show that σ -conjugating x by certain very simple elements of $G(L)$ corresponds to adding “appendages” to the ends of Γ . One can then understand how the addition of such appendages affects relative position. Some of the “appendage” methods used are stated in generality in Section 3.5.

The second method of producing subsets of solution sets is lengthy and unappealing, but applies for $b \neq 1$. The basic idea is as follows. In Section 3.1.1, we produce a gallery Γ_x between xC_M and $b\sigma(x)C_M$. We compute the folding results of $x^{-1}\Gamma_x$ to get possible candidates for the relative position of x and $b\sigma(x)$. In Section 3.1.6,

we discover which of these candidates can actually occur by defining some invariants which control how $x^{-1}\Gamma_x$ actually folds (i.e., these invariants control which of the folding results of $x^{-1}\Gamma_x$ is the actual value of the last chamber of $\rho(x^{-1}\Gamma_x)$). We then show that one can choose x such that these invariants take any pre-specified value. One of the complications of this method is that one must proceed somewhat differently for each of the three cases $b = 1$, b “degenerate” but $b \neq 1$, and b “non-degenerate.” The “non-degenerate” b are exactly

$$b = \begin{pmatrix} \pi^\alpha & 0 & 0 \\ 0 & \pi^\beta & 0 \\ 0 & 0 & \pi^\gamma \end{pmatrix}$$

with $\alpha + \beta + \gamma = 0$, $\alpha > \beta > \gamma$.

CHAPTER 2

GROUPS OF SEMISIMPLE RANK 1

If the split connected reductive group G is of semisimple rank 1, then it is possible to obtain a complete characterization of which $X_w(b\sigma)$ are non-empty. Some of the methods which will be used on higher rank groups are similar to those used here. Considering the rank 1 case first allows one to develop a good intuition about the geometric nature of the problem before proceeding to the more complicated rank 2 and higher rank cases.

Section 2.1 of this chapter will address the specific group SL_2 , and Section 2.2 will adapt the SL_2 solution to other, non-simply connected groups of semisimple rank 1.

Everything in this chapter was previously known to Kottwitz and Rapoport.

2.1 SL_2

Results from a paper of Kottwitz [6] allow us to list the following elements of SL_2 as a complete collection of representatives of the σ -conjugacy classes of SL_2 :

$$\left\{ \begin{pmatrix} \pi^n & 0 \\ 0 & \pi^{-n} \end{pmatrix} : n \in \mathbb{Z}, n \geq 0 \right\}.$$

Let $\rho : \mathcal{B}_\infty \rightarrow A_M$ be the retraction of $\mathcal{B}_\infty$ onto A_M relative to C_M .

In Section 2.1.1, we answer the question of which w have non-empty $X_w(b\sigma)$ for b the identity matrix. In Section 2.1.2, we consider the cases where b is one of the above non-identity representatives. In Section 2.1.3, we prove that the results obtained in Sections 2.1.1 and 2.1.2 are invariant under a certain $\mathbb{Z}/2$ -action. The proof is independent of the results from Sections 2.1.1 and 2.1.2. Section 2.1.4 describes the relationship between $X_w(1\sigma)$ and $X_w(b\sigma)$ for $b \neq 1$.

2.1.1 Identity σ -conjugacy Class

As mentioned previously, to find out which w have non-empty $X_w(b\sigma)$, it suffices to determine the set $\{inv(1, x^{-1}b\sigma(x)) : x \in SL_2(L)\}$. However, it is easy to see that $inv(1, x^{-1}b\sigma(x)) = \rho(x^{-1}b\sigma(x)C_M)$, where again it is understood that we are making use of the correspondence between $\tilde{W}$ and the set of chambers in A_M . So we seek to describe the set $\{\rho(x^{-1}b\sigma(x)C_M) : x \in SL_2(L)\}$.

Let D be any chamber in $\mathcal{B}_\infty$.

Definition 2.1.1. Let the distance $\tilde{d}_1(D)$ between D and $\mathcal{B}_1$ be $n - 1$, where n is the length of the shortest gallery containing D and some chamber E in $\mathcal{B}_1$.




Figure 2.1: Chamber labels




Figure 2.2: Alternate chamber labels

Note that D is in $\mathcal{B}_1$ if and only if $\tilde{d}_1(D) = 0$.

Proposition 2.1.1. If $\tilde{d}_1(D) > 0$ then there is a minimal gallery between D and $\sigma(D)$ of length $2\tilde{d}_1(D)$.

Proof. Let G be a gallery of minimal length containing D and some chamber E in $\mathcal{B}_1$. Let $E = G_1, G_2, G_3, \dots, G_n = D$ be the successive chambers of G . Then G_2 is not contained in $\mathcal{B}_1$, or G would not be of minimal length. So the gallery $G_n, G_{n-1}, \dots, G_2, \sigma(G_2), \dots, \sigma(G_{n-1}), \sigma(G_n)$ is non-stuttering and therefore minimal. $\square$

Now let $D = xC_M$. By the nature of the SL_2 action on the building, x can be chosen such that $\tilde{d}_1(xC_M)$ is any non-negative integer. Therefore, x can be chosen such that the minimal gallery between xC_M and $\sigma(xC_M) = \sigma(x)C_M$ is any even integer greater than or equal to 2. In the case where $\tilde{d}_1(xC_M) = 0$ (i.e., xC_M is contained in $\mathcal{B}_1$), we have $xC_M = \sigma(xC_M)$.

We are interested in $\rho(x^{-1}\sigma(x)C_M)$, so the minimal gallery between xC_M and $\sigma(x)C_M$ constructed above is useful. We will need the following terminology to describe the set $\{\rho(x^{-1}\sigma(x)C_M) : x \in SL_2(L)\}$.

Definition 2.1.2. Let C_M^i denote the chamber in A_M as described in Figure 2.1, where $C_M^0 = C_M$.

Note that the choice between this labeling and the one in Figure 2.2 is arbitrary, but we choose the former.

Proposition 2.1.2. The $w \in \tilde{W}$ with non-empty $X_w(1\sigma)$ are C_M and C_M^i for i odd.

Proof. First, note that $C_M = \rho(x^{-1}\sigma(x)C_M)$ for x such that xC_M is contained completely in $\mathcal{B}_1$. Further, since x can be chosen such that the minimal gallery between xC_M and $\sigma(xC_M)$ is of any even integral length, we see that for each odd




Figure 2.3: Proof of Proposition 2.1.2

i , either C_M^i or C_M^{-i} is obtained. We must show that in fact both are obtained. For this, note that if $\rho(x^{-1}\sigma(x)C_M) = C_M^i$ then $\rho(y^{-1}\sigma(y)C_M) = C_M^{-i}$ for $y = gxq^{-1}$ where

$$g = \begin{pmatrix} \pi & 0 \\ 0 & 1 \end{pmatrix} \quad ; \quad q = \begin{pmatrix} 0 & -\pi \\ 1 & 0 \end{pmatrix}.$$

See Figure 2.3. In this figure, the vertex types are labeled. The key point is that the vertices on the chambers xC_M are of types opposite to the corresponding vertices on $gxq^{-1}C_M$. Note that we still have $\det(y) = 1$. See Section 2.1.3 for a reminder of how q acts on C_M and A_M . $\square$

2.1.2 Non-identity σ -conjugacy Classes

Again, let D be any chamber in $\mathcal{B}_\infty$. Let $b = \begin{pmatrix} \pi^s & 0 \\ 0 & \pi^{-s} \end{pmatrix}$, $s > 0$. We will enumerate the w for which $X_w(b\sigma)$ is non-empty.

Definition 2.1.3. The distance $\tilde{d}_{A_M}(D)$ between D and A_M is $n - 1$, where n is the length of the shortest gallery containing D and some chamber E in A_M .

Note that D is in A_M if and only if $\tilde{d}_{A_M}(D) = 0$.

Proposition 2.1.3. If $\tilde{d}_{A_M}(D) > 0$ then there exists a minimal gallery between D and $b\sigma(D)$ of length $2s + 2\tilde{d}_{A_M}(D)$.

Proof. Let G be a gallery of minimal length containing D and some chamber E in A_M . Let $E = G_1, G_2, G_3, \dots, G_n = D$ be the successive chambers of G . Here $n = \tilde{d}_{A_M}(D) + 1$. Then G_2 is not contained in A_M , or G would not be of minimal length. Let v be the vertex that E and G_2 share. There is a unique minimal gallery contained in A_M between v and $b\sigma(v) = bv$. Call this gallery H , and its chambers $H_1, H_2, \dots, H_{2s}$, where v is the vertex of H_1 that is not also a vertex of H_2 . Consider the gallery $G_n, G_{n-1}, \dots, G_2, H_1, H_2, \dots, H_{2s}, b\sigma(G_2), b\sigma(G_3), \dots, b\sigma(G_n)$. This is non-stuttering and therefore minimal. It has the desired length. $\square$

Now consider $D = xC_M$. Clearly, x can be chosen such that $\tilde{d}_{A_M}(xC_M)$ is any non-negative integer. Therefore, x can be chosen such that the minimal gallery between xC_M and $b\sigma(xC_M)$ is of length $2s + 2n$ for any $n \geq 1$. If xC_M is contained in A_M then the minimal gallery between xC_M and $b\sigma(xC_M)$ is of length $2s + 1$.

Proposition 2.1.4. The $w \in \tilde{W}$ with non-empty $X_w(b\sigma)$ are $C_M^{2s}, C_M^{-2s}, C_M^{2s+i}$ and C_M^{-2s-i} for $i \geq 1$ any odd integer.

Proof. Similar to Proposition 2.1.2 in Section 2.1.1. $\square$

Figure 2.4 summarizes results from this and the previous section.


2.1.3 Symmetry Under a $\mathbb{Z}/2$ -action

Let $q = \begin{pmatrix} 0 & -\pi \\ 1 & 0 \end{pmatrix} \in GL_2(L)$. Since $\mathcal{B}_\infty$ is also the building for $GL_2(L)$, we can ask how q acts on A_M . The main apartment can be represented using lattice classes as in Figure 2.5. The matrix q takes

$$\begin{aligned} \mathcal{O} \oplus \mathcal{O} &\rightarrow \pi\mathcal{O} \oplus \mathcal{O} \\ \pi\mathcal{O} \oplus \mathcal{O} &\rightarrow \mathcal{O} \oplus \mathcal{O} \\ \pi^{-1}\mathcal{O} \oplus \mathcal{O} &\rightarrow \pi^2\mathcal{O} \oplus \mathcal{O} \\ \pi^2\mathcal{O} \oplus \mathcal{O} &\rightarrow \pi^{-1}\mathcal{O} \oplus \mathcal{O} \end{aligned}$$

and therefore represents a flip about the midpoint of C_M . One can see by looking at the results of the previous two sections that $\{w \in \tilde{W} : X_w(b\sigma) \neq \emptyset\}$ is invariant under this q action. It is instructive to develop a proof of this fact that is independent of the previous sections.

Since inv is a map from $SL_2(L)/I \times SL_2(L)/I \rightarrow \tilde{W}$, and since $SL_2(L)/I$ is in one-to-one correspondence with elements of the set of chambers in $\mathcal{B}_\infty$, we can view inv as a map that takes two chambers in $\mathcal{B}_\infty$ as arguments. We will use both this and the standard method of referring to inv throughout the rest of the paper.

Figure 2.4: Summary of SL_2 results




Figure 2.5: Lattice labels for vertices

We know that $\text{inv}(xC_M, b\sigma(x)C_M) = \text{inv}(gxC_M, gb\sigma(x)C_M)$ for any $g \in SL_2(L)$. If $g \in SL_2(F)$ is diagonal then $\text{inv}(xC_M, b\sigma(x)C_M) = \text{inv}(gxC_M, gb\sigma(x)C_M) = \text{inv}(gxC_M, b\sigma(gx)C_M)$. In the case that $g \in GL_2(F)$ is diagonal, $\text{inv}(xC_M, b\sigma(x)C_M)$ is not necessarily equal to $\text{inv}(gxC_M, gb\sigma(x)C_M)$, but $\text{inv}(gxC_M, gb\sigma(x)C_M) = \text{inv}(gxC_M, b\sigma(gx)C_M)$. If y is such that $yC_M = gxC_M$, and $y \in SL_2(L)$, then $\text{inv}(gxC_M, b\sigma(gx)C_M) = \text{inv}(yC_M, b\sigma(y)C_M)$ is a new $w \in \tilde{W}$ such that $X_w(b\sigma) \neq \emptyset$. We used the fact that if $yC_M = gxC_M$ then $b\sigma(gx)C_M = b\sigma(gxC_M) = b\sigma(yC_M) = b\sigma(y)C_M$.

Proposition 2.1.5. We can choose g a diagonal element of $GL_2(F)$ such that if $\text{inv}(xC_M, b\sigma(x)C_M) = C_M^i$, then $\text{inv}(gxC_M, b\sigma(gx)C_M) = C_M^{-i}$.

Proof. Let $g = \begin{pmatrix} \pi & 0 \\ 0 & 1 \end{pmatrix}$. We see that $\text{inv}(gxC_M, b\sigma(gx)C_M) = \rho(s^{-1}b\sigma(gx)C_M)$, where $s \in SL_2(L)$ is chosen such that $s^{-1}gxC_M = C_M$. We may choose s such that $s^{-1} = qx^{-1}g^{-1}$. So $\rho(s^{-1}b\sigma(gx)C_M) = \rho(qx^{-1}g^{-1}bg\sigma(x)C_M) = \rho(qx^{-1}b\sigma(x)C_M)$, and if $\text{inv}(xC_M, b\sigma(x)C_M) = C_M^i$, then $\rho(qx^{-1}b\sigma(x)C_M) = C_M^{-i}$. $\square$

We will later adapt this proof to give a similar result for some groups of higher rank.


2.1.4 Relationship Between $X_w(1\sigma)$ and $X_w(b\sigma)$

In this section we explore the relationship between the set of w such that $X_w(1\sigma)$ is non-empty and the set of w such that $X_w(b\sigma)$ is non-empty, for some $b \neq 1$.

Proposition 2.1.6. If w is in the positive Weyl chamber (i.e., $w = C_M^i$ for $i \geq 0$), then $X_w(1\sigma)$ is non-empty if and only if $X_{bw}(b\sigma)$ is non-empty. If w is in the negative Weyl chamber (i.e., $w = C_M^i$ for $i < 0$), then $X_w(1\sigma)$ is non-empty if and only if $X_{b^{-1}w}(b\sigma)$ is non-empty.

Proof. We start by remarking that this is clear from the results of Section 2.1.1 and Section 2.1.2. But we produce an *a priori* proof that may be generalizable in some respects to higher rank groups.

Assume that w is in the positive Weyl chamber. If $X_w(1\sigma)$ is non-empty then let x be such that $\text{inv}(x, \sigma(x)) = w$. Let A_1 be an apartment containing xC_M and C_M . Let G be the minimal gallery (in A_1) between C_M and xC_M , and let

Figure 2.6: For w in the positive Weyl chamber

$C_M = G_1, G_2, \dots, G_s = xC_M$ be the chambers of G . Let i be maximal such that G_i is contained in $\mathcal{B}_1$. We could have $G_i = xC_M$ in the case that $w = C_M$. Let A_2 be an apartment containing C_M and G_i . If $i \neq s$ then we also require that A_2 not contain G_{i+1} . Let $g \in SL_2(L)$ be such that $gC_M = C_M$ and g sends A_2 to A_M . To see that $\text{inv}(gx C_M, b\sigma(gx)C_M) = bw$, consider Figure 2.6 in which $b = \begin{pmatrix} \pi^s & 0 \\ 0 & \pi^{-s} \end{pmatrix}$.

The minimal gallery E_2 between $gx C_M$ and $b\sigma(gx)C_M$ is $2s$ chambers longer than the minimal gallery E_1 between xC_M and $\sigma(x)C_M$, and the types of the vertices on the chamber $gx C_M$ are the same as those of the corresponding vertices of xC_M . So if $\rho(x^{-1}E_1)$ is the minimal gallery in A_M between C_M and w then $\rho((gx)^{-1}E_2)$ is the minimal gallery in A_M between C_M and bw . To get the other implication, use g^{-1} .

Now assume that w is in the negative Weyl chamber. Let G, G_i, A_2 and g be as before. In this situation, Figure 2.6 becomes Figure 2.7. As before, the gallery E_2 between $gx C_M$ and $b\sigma(gx)C_M$ is $2s$ chambers longer than the minimal gallery E_1 between xC_M and $\sigma(x)C_M$, and the types of the vertices on the chamber $gx C_M$ are the same as those of the corresponding vertices of xC_M . So if $\rho(x^{-1}E_1)$ is the minimal gallery in A_M between C_M and w then $\rho((gx)^{-1}E_2)$ is the minimal gallery in A_M that begins at C_M and goes in the same direction as $\rho(x^{-1}E_1)$, but that is $2s$ chambers longer. In the current case, the final chamber of this gallery is $\tilde{b}w$.

As before, to get the other implication, use g^{-1} . □




Figure 2.7: For w in the negative Weyl chamber

2.2 GL_2 and PGL_2

The situation is slightly different for GL_2 . First, we must understand differences in setup. For $GL_2(L)$ it is possible to send C_M to itself without fixing it pointwise. The matrix $m = \begin{pmatrix} 0 & \pi \\ 1 & 0 \end{pmatrix}$ does this, for example. Another way of saying this is that contrary to the SL_2 situation, the Iwahori associated to C_M is no longer the set $\{g : gC_M = C_M\}$.

We still have $inv : GL_2(L)/I \times GL_2(L)/I \rightarrow I \backslash GL_2(L)/I \simeq \tilde{W}$, but now $\tilde{W} \simeq W_a \rtimes \mathbb{Z}$, where W_a , the affine Weyl group, is in one-to-one correspondence with the chambers of A_M (recall that for SL_2 , we had $W_a \simeq \tilde{W}$). We therefore have $inv : GL_2(L)/I \times GL_2(L)/I \rightarrow W_a \rtimes \mathbb{Z}$. This map is given by $inv(x, y) = (\rho(x^{-1}yC_M), v(\det(x^{-1}y)))$, where v is the valuation.

The σ -conjugacy classes of $GL_2(L)$ are

$$\left\{ \begin{pmatrix} \pi^\alpha & 0 \\ 0 & \pi^\beta \end{pmatrix} : \alpha \geq \beta \right\} \cup \left\{ \begin{pmatrix} 0 & \pi \\ 1 & 0 \end{pmatrix}^\alpha : \alpha \in \mathbb{Z}, \alpha \text{ odd} \right\} = R.$$


This follows from a result of Kottwitz [6].



For a given σ -conjugacy class $\{x^{-1}b\sigma(x) : x \in GL_2(L)\}$, where $b \in R$, we are interested in which $(w, n) \in W_a \rtimes \mathbb{Z}$ have non-empty $X_{(w,n)}(b\sigma)$. To solve this, we will describe the set $\{(\rho(x^{-1}b\sigma(x)C_M), v(\det(x^{-1}b\sigma(x)))) \in W_a \rtimes \mathbb{Z} : x \in GL_2(L)\}$ for any fixed b in the above set R .

First, note that $v(\det(x^{-1}b\sigma(x))) = v(\det(b))$, so the second component is fixed for fixed b . The possible values of the first component can be determined using a process that is similar to that in Sections 2.1.1 and 2.1.2. The difference is that b can now take on more values. Also, it is *a priori* possible (although we will see that it does not occur) for $\{\rho(x^{-1}b\sigma(x)C_M) : x \in GL_2(L)\}$ to be bigger than $\{\rho(x^{-1}b\sigma(x)C_M) : x \in SL_2(L)\}$, since $GL_2(L)$ acts on $\mathcal{B}_\infty$ in ways that $SL_2(L)$ does not.

If $b = \begin{pmatrix} \pi^\alpha & 0 \\ 0 & \pi^\beta \end{pmatrix}$ then b shifts A_M to the right by $\alpha - \beta$ units. Using reasoning similar to that in Section 2.1.2, for these b we see that $\rho(x^{-1}b\sigma(x)C_M)$ could be any of the chambers $C_M^{\pm(\alpha-\beta)}$ or $C_M^{\pm(\alpha-\beta+i)}$, for all i odd, $i \geq 1$. Figure 2.8 summarizes these results.

Now we consider $b = \begin{pmatrix} 0 & \pi \\ 1 & 0 \end{pmatrix}^\alpha$, $\alpha \in \mathbb{Z}, \alpha$ odd. The matrix $m = \begin{pmatrix} 0 & \pi \\ 1 & 0 \end{pmatrix}$ flips A_M about the center of C_M . Therefore so does the matrix b . Let G be the minimal gallery between C_M and xC_M . Then $b\sigma(G)$ is the minimal gallery between C_M and $b\sigma(x)C_M$. Let $C_M = G_1, G_2, \dots, G_s = xC_M$ be the chambers of G . Then $G_s, G_{s-1}, \dots, G_1, b\sigma(G_2), b\sigma(G_3), \dots, b\sigma(G_s)$ is the unique minimal gallery between xC_M and $b\sigma(x)C_M$. Call this gallery Γ_x . It has length $2s - 1$.

Figure 2.8: Summary of some results for GL_2

Figure 2.9: Computations for some σ -conjugacy classes for GL_2 Figure 2.10: Summary of some more results for GL_2

By choosing x appropriately, we can arrange for s to be any integer ≥ 1 . Further, if $\rho(x^{-1}\Gamma_x)$ has final chamber C_M^i , then $\rho(y^{-1}\Gamma_y)$ has final chamber C_M^{-i} , where $y = xm$. Therefore, the possible values for $\rho(x^{-1}b\sigma(x)C_M)$ are C_M^i for any i even. This is independent of which odd $\alpha \in \mathbb{Z}$ we have chosen. Figure 2.9 makes the calculations more clear, and Figure 2.10 summarizes the possible values of the first component of inv . The second component is always $v(\det(x^{-1}b\sigma(x))) = v(\det(b))$.

For PGL_2 , $\tilde{W} \cong W_a \ltimes \mathbb{Z}/2$, and $inv(x, y) = (\rho(x^{-1}yC_M), v(\det(x^{-1}y)))$, where in PGL_2 , $v(\det(x^{-1}y))$ is only determined mod 2. The results for PGL_2 are the same as those for GL_2 , but second-component values are computed mod 2.

2.3 Reorganization of Results

So far, we have stated all results by specifying, for each b , which w have non-empty $X_w(b\sigma)$, or which (w, n) have non-empty $X_{(w, n)}(b\sigma)$. We reword these results by saying, for each w , which b have non-empty $X_w(b\sigma)$, and for each (w, n) , which b have non-empty $X_{(w, n)}(b\sigma)$.

We first reword results for SL_2 . If $b = \begin{pmatrix} \pi^s & 0 \\ 0 & \pi^{-s} \end{pmatrix}$, then let Σ_s be the σ -conjugacy class of b . Let D_M^i denote the double- I -coset corresponding to the chamber C_M^i . Then one can see that $D_M^{\pm 2n} \subseteq \Sigma_n$ for $n \in \mathbb{Z}$, $n \geq 0$. One can also see that for $n \geq 0$,

i	j
0	0
2	1
4	2
6	3
$\vdots$	$\vdots$

Table 2.1: $D_M^{\pm i}$ intersects exactly these Σ_j non-trivially for i even

i	j
1	0
3	0, 1
5	0, 1, 2
7	0, 1, 2, 3
$\vdots$	$\vdots$

Table 2.2: $D_M^{\pm i}$ intersects exactly these Σ_j non-trivially for i odd

$n \in \mathbb{Z}$, $D_M^{\pm(2n+1)}$ intersects the σ -conjugacy classes $\Sigma_0, \Sigma_1, \dots, \Sigma_n$, and no others. These results are summarized in Tables 2.1 and 2.2.

In particular, D_M^{2n} for $n \in \mathbb{Z}$ and $D_M^{\pm 1}$ are each completely contained in some σ -conjugacy class. The cosets $D_M^{\pm(2n+1)}$ for $n \in \mathbb{Z}$, $n \geq 0$ are spread over increasingly many σ -conjugacy classes as n gets larger.

We now reword results for GL_2 . Let $\Sigma_{\alpha, \beta}$ be the σ -conjugacy class of $\begin{pmatrix} \pi^\alpha & 0 \\ 0 & \pi^\beta \end{pmatrix}$, and let Σ_α be the σ -conjugacy class of $\begin{pmatrix} 0 & \pi \\ 1 & 0 \end{pmatrix}^\alpha$. Let D_M^i denote the element of W_a corresponding to the chamber C_M^i in A_M , and let (D_M^i, j) be an element of $\tilde{W} \cong W_a \rtimes \mathbb{Z}$. We also denote the corresponding double- I -coset by (D_M^i, j) . Then one can see that $(D_M^i, j) \subseteq \Sigma_{\frac{j+i}{2}, \frac{j-i}{2}}$ if $i \equiv j \pmod{2}$. If i is even and j is odd, then (D_M^i, j) intersects Σ_j and $\Sigma_{\frac{j+k}{2}, \frac{j-k}{2}}$ for $1 \leq k \leq i-1$ an odd integer. It does not intersect non-trivially with any other σ -conjugacy classes. Finally, if i is odd and j is even, then (D_M^i, j) intersects $\Sigma_{\frac{j+k}{2}, \frac{j-k}{2}}$ for $0 \leq k \leq i-1$ an even integer. It does not intersect non-trivially with any other σ -conjugacy classes. These results are summarized in Table 2.3.

i	j odd	j even
0	Σ_j	$\Sigma_{\frac{j}{2}, \frac{j}{2}}$
± 1	$\Sigma_{\frac{j+1}{2}, \frac{j-1}{2}}$	$\Sigma_{\frac{j}{2}, \frac{j}{2}}$
± 2	$\Sigma_j ; \Sigma_{\frac{j+1}{2}, \frac{j-1}{2}}$	$\Sigma_{\frac{j+2}{2}, \frac{j-2}{2}}$
± 3	$\Sigma_{\frac{j+3}{2}, \frac{j-3}{2}}$	$\Sigma_{\frac{j}{2}, \frac{j}{2}} ; \Sigma_{\frac{j+2}{2}, \frac{j-2}{2}}$
± 4	$\Sigma_j ; \Sigma_{\frac{j+1}{2}, \frac{j-1}{2}} ; \Sigma_{\frac{j+3}{2}, \frac{j-3}{2}}$	$\Sigma_{\frac{j+4}{2}, \frac{j-4}{2}}$
± 5	$\Sigma_{\frac{j+5}{2}, \frac{j-5}{2}}$	$\Sigma_{\frac{j}{2}, \frac{j}{2}} ; \Sigma_{\frac{j+2}{2}, \frac{j-2}{2}} ; \Sigma_{\frac{j+4}{2}, \frac{j-4}{2}}$
± 6	$\Sigma_j ; \Sigma_{\frac{j+1}{2}, \frac{j-1}{2}} ; \Sigma_{\frac{j+3}{2}, \frac{j-3}{2}} ; \Sigma_{\frac{j+5}{2}, \frac{j-5}{2}}$	$\Sigma_{\frac{j+6}{2}, \frac{j-6}{2}}$
$\vdots$	$\vdots$	$\vdots$

Table 2.3: σ -conjugacy classes that intersect (D_M^i, j) non-trivially

CHAPTER 3

GROUPS OF SEMISIMPLE RANK 2

In Section 3.1 we will show for b an element of a certain collection of σ -conjugacy class representatives of $SL_3(L)$, exactly which double- I -cosets intersect $\{x^{-1}b\sigma(x) : x \in SL_3(L)\}$ non-trivially. In Section 3.2, we will show that these results can also be made to give complete information of the same kind for certain σ -conjugacy classes of $GL_3(L)$ and $PGL_3(L)$. Section 3.3 attempts to apply the same methods to Sp_4 . Certain results are conjectural in this setting because some of the lengthy computations done for SL_3 have not been done in entirety for Sp_4 . Section 3.4 discusses how the methods of Section 3.1 could be applied to other rank 2 groups. Section 3.5 gives some invariance properties of the set $\{inv(x, b\sigma(x)) : x \in G(L)\}$ that hold for any simply-connected group G , and any σ -conjugacy class b . These could be applied to higher rank groups, for instance. Some of the methods of Section 3.1 could also be applied to SL_n or Sp_{2n} . A comment to this effect is made when this is the case.

3.1 SL_3

In this section we enumerate the non-empty $X_w(b\sigma)$ for SL_3 , where the letter b will always refer to an element of $SL_3(L)$ of the form

$$\begin{pmatrix} \pi^\alpha & 0 & 0 \\ 0 & \pi^\beta & 0 \\ 0 & 0 & \pi^{-\alpha-\beta} \end{pmatrix},$$

with $\alpha \geq \beta \geq \alpha - \beta$. These are representatives of distinct σ -conjugacy classes of $SL_3(L)$. However, not every σ -conjugacy class is represented by one of these elements [6].

The collection of chambers in A_M that correspond to w with non-empty $X_w(b\sigma)$ is computed in the following way. In Section 3.1.1, we give some necessary definitions. In Section 3.1.2, we use geometric methods to produce a superset of the desired collection of chambers. In Section 3.1.4 we use algebraic methods that produce a subset of the desired collection for $b = 1$. We enlarge this subset in Section 3.1.5. This enlargement will be equal to the superset from Section 3.1.2, and therefore is the solution set of chambers. In Section 3.1.6 we give a geometric method of arriving at the subset of Section 3.1.5 that also applies for $b \neq 1$. In Section 3.1.7, we prove a symmetry result analogous to that done for SL_2 in Section 2.1.3. In Section 3.1.3, we develop a relationship between the collection of w with non-empty $X_w(1\sigma)$ and

the collection of w with non-empty $X_w(b\sigma)$. This is done by illuminating an efficient method of computing $\{inv(x, b\sigma(x)) : x \in SL_3(L)\}$ that works for any of the b listed above.

3.1.1 Standard Minimal Galleries and Composite Galleries

Our goal is to determine which $X_w(b\sigma)$ are non-empty. The technique that led to success in the rank 1 case involved considering, for every chamber $E \subseteq \mathcal{B}_\infty$, a gallery Γ_E connecting E to $b\sigma(E)$. We used the unique minimal gallery between these two chambers. We will use a similar process for SL_3 . However, there is no longer a unique minimal gallery Γ_E connecting E and $b\sigma(E)$. In this section we will specify a choice of gallery, which we will call the *composite gallery*. The composite gallery is usually not minimal.

First define the three *primary directions* D_1, D_2, D_3 and the three *secondary directions* d_1, d_2, d_3 in the main apartment A_M for SL_3 as marked by arrows in Figure 3.1. Given a chamber $E \subseteq A_M$, we define the *standard minimal gallery* (SMG) between C_M and E as follows. If E is in one of the corridors marked $c_1, \dots, c_6$ in Figure 3.1, then the SMG is the unique minimal gallery from C_M to E . If E is in region R_i , proceed first in the direction D_i , then in the direction d_i . If E is in region r_i , proceed first in direction D_i , then in direction d_j , where $j \equiv i - 1 \pmod{3}$.

The following lemma allows us to define the SMG between C_M and E for any chamber $E \subseteq \mathcal{B}_\infty$ in a natural way. As in Chapter 2, ρ is the retraction from $\mathcal{B}_\infty$ to A_M centered at C_M .

Lemma 3.1.1. There is a unique gallery G_E between C_M and E which is minimal, and such that $\rho(G_E)$ is the SMG between C_M and $\rho(E)$.

Proof. Let A_1 be an apartment containing C_M and E . Let $g_1 \in SL_3(L)$ be such that $g_1 A_1 = A_M$ and $g_1 C_M = C_M$. If $G_{\rho(E)}$ is the SMG between C_M and $\rho(E)$, then let $G_E = g_1^{-1} G_{\rho(E)}$. This proves existence.

To prove uniqueness, note that if $\tilde{G}_E$ is another minimal gallery from C_M to E , then $\tilde{G}_E \subseteq A_1$ (see Definition 3.1.6 and Theorem 3.1.6 in Section 3.1.6). If $\rho(\tilde{G}_E) = G_{\rho(E)}$ then $g_1^{-1} G_{\rho(E)} = \tilde{G}_E$. But $g_1^{-1} G_{\rho(E)} = G_E$. $\square$

To obtain a composite gallery, Γ_E , from the SMG between C_M and E , proceed as follows.

Case 1: $b = 1$. Let Γ_E^1 be the gallery composed of the chambers of the SMG between C_M and E that are not in $\mathcal{B}_1$. Let $\Gamma_E^3 = \sigma(\Gamma_E^1)$. Then $\Gamma_E = \Gamma_E^1 \cup \Gamma_E^3$. We will call the edge e between Γ_E^1 and Γ_E^3 the *edge of departure of the SMG between C_M and E from $\mathcal{B}_1$* .

Case 2: $b \neq 1$. Let Γ_E^1 be the gallery composed of the chambers of the SMG between C_M and E that are not in A_M . Let $\Gamma_E^3 = b\sigma(\Gamma_E^1)$. Let e be the unique edge in Γ_E^1




Figure 3.1: Primary and secondary directions

that is contained in A_M . Let Γ_E^2 be any minimal gallery connecting e and $b\sigma(e) = be$. Then $\Gamma_E = \Gamma_E^1 \cup \Gamma_E^2 \cup \Gamma_E^3$. We call e the *edge of departure of the SMG between C_M and D from A_M* . The term *edge of departure* without further modification will be used to refer either to an edge of departure from A_M or to an edge of departure from $\mathcal{B}_1$.

We will also need the following definitions.

Definition 3.1.1. If E_1 and E_2 are two chambers in $\mathcal{B}_\infty$ that share an edge e , then we say the *transition type* from E_1 to E_2 is the type of the two vertices not in e .

Definition 3.1.2. If G is a non-stuttering gallery in $\mathcal{B}_\infty$ consisting of $G_0, \dots, G_s$, and if the transition type from G_{i-1} to G_i is t_i for $1 \leq i \leq s$, then we say that G has *type* $t_1, \dots, t_s$.

Note that if $G \subseteq A_M$, then G_0 and $t_1, \dots, t_s$ determine G . It is also possible to specify a gallery type by giving a picture of a non-stuttering gallery in A_M .

3.1.2 A Superset of the Solution Set

Just as in the SL_2 case, SL_3 is simply connected. So given b , an answer to the question of which $w \in \tilde{W}$ have non-empty $X_w(b\sigma)$ can be given by specifying the corresponding chambers in A_M . As such, the answer we are looking for is the set of chambers $S = \{\rho(x^{-1}b\sigma(x)C_M) : x \in SL_3(L)\}$. In the SL_2 case, for every $x \in SL_2(L)$, we had a unique minimal gallery Γ_x connecting xC_M and $\sigma(x)C_M$. Since this gallery was minimal, we were able to determine $\rho(x^{-1}b\sigma(x)C_M)$ just by folding $x^{-1}\Gamma_x$ from C_M down into A_M in the unique possible way. In the SL_3 case, we let Γ_x instead be the composite gallery Γ_E (for $E = xC_M$) constructed in the previous section. Since this composite gallery is not necessarily minimal, one can not determine $\rho(x^{-1}b\sigma(x)C_M)$ from its type and starting point alone. One can, however, get a set of possible chambers S_x for $\rho(x^{-1}b\sigma(x)C_M)$ by enumerating all the possible foldings of a gallery of the same type as $x^{-1}\Gamma_x$, and putting the final chamber of each into S_x . The set $S_1 = \cup_{x \in SL_3(L)} S_x$ contains the set S in which we are interested.

We include an example of this computational process. Let

$$b = \begin{pmatrix} \pi^3 & 0 & 0 \\ 0 & \pi^{-1} & 0 \\ 0 & 0 & \pi^{-2} \end{pmatrix},$$

and let x be such that the SMG connecting C_M to xC_M has type and edge of departure from A_M as indicated in Figure 3.2. Then the resulting composite gallery Γ_x has the same type as the gallery pictured in Figure 3.3. Therefore, $\rho(x^{-1}b\sigma(x)C_M)$ must be one of the chambers marked on Figure 3.4. These results are achieved by computing all possible values of $\rho(E)$, where E is the last chamber of some gallery that begins at C_M and that has the same type as Γ_x (pictured in Figure 3.3).




Figure 3.2: Example SMG type and edge of departure




Figure 3.3: Composite gallery example





Figure 3.4: Results from the SMG in Figure 3.2

The apparent problem with this process is that it seems to be an infinite computation. The set S_1 is a union of the S_x , where $x \in SL_3(L)$ is arbitrary. Of course, we need only consider one representative x from each coset $SL_3(L)/I$, but there are still infinitely many such cosets. We can further optimize using the fact that only the *type* of $x^{-1}\Gamma_x$ is important for computing S_x . This type is determined by the type of the SMG between C_M and xC_M , and the departure edge of this SMG from A_M (or from $\mathcal{B}_1$ if $b = 1$). We call a pair consisting of an SMG type and a departure edge a *type-edge pair*. The set we are trying to compute is $S_1 = \cup S_{(t,e)}$, where the union is over all type-edge pairs (t,e) , and where $S_{(t,e)} = S_x$ for some $x \in SL_3(L)$ such that the SMG from C_M to xC_M has type t and departure edge e .

The benefit of this point of view is that the $S_{(t,e)}$ can be separated into finitely many infinite classes, each of which can be computed all at once. We give an example of two of these infinite classes, I_1 and I_2 , for b as above. The SMG types in I_1 are those represented by the SMGs of the chambers in the region R_2 in Figure 3.1. The edges of departure from A_M that we will consider are the horizontal ones. The SMG types in I_2 are those represented by the SMGs of the chambers in corridor c_4 . The pairs $(t,e) \in I_2$ will have arbitrary e .

We consider I_1 first. As in Case 2 of the definition of the composite gallery, let Γ_E^1 be the gallery composed of the chambers of the SMG in question that are not in A_M (i.e., those after the edge of departure). So I_1 gives rise to the Γ_E^1 in Figure 3.5.

Let W be the finite Weyl group, and let $w \in W$, $a \in T(F)$ be such that $awC_M =$

Figure 3.5: Γ_E^1 possibilities for I_1

$\rho(E)$. We break I_1 into six sub-classes according to w . Let

$$f = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

Case 1: $w = f$. We get a composite gallery of the type shown in Figure 3.6. If the *turning edge* is labeled in Figure 3.5, then we consider in Figure 3.7 the instances where Γ_E^1 has 3 chambers after the turning edge. We consider in Figure 3.8 the instances where Γ_E^1 has 5 chambers after the turning edge. We consider on Figure 3.9 the instances where Γ_E^1 has 7 chambers after the turning edge. After doing these computations, it is easy to see what the situation is for the instances where Γ_E^1 has $2n + 1$ chambers after the turning edge, for $n \geq 1$. The results for these n are put together in Figure 3.10.

Case 2: $w = 1$. We get a composite gallery of the type shown in Figure 3.11. Figures 3.12, 3.13, and 3.14 have instances where Γ_E^1 has 2, 4, and 6 chambers after the turning edge, respectively. Figure 3.15 has the amalgamated results for $2n$ chambers after the turning edge for all $n \geq 1$.

It is easy to see that the results for the cases in which w is some other order 2 element of W are just rotations of the results for case 1 by 120° and 240° about the center point of the chamber C_M . The results for the cases in which w is some other order 3 element of W are rotations of the case 2 results by 120° and 240° about the center point of C_M .

We break I_2 into the same six sub-classes.

Case 1: $w = 1$. We get a composite gallery of one of the two types shown in Figures 3.16 and 3.17, where D is the last chamber in the SMG from C_M to xC_M that is contained in A_M . We consider in Figure 3.18 the instances where Γ_E^1 has 0, 1 or 2 chambers after the departure edge, in Figure 3.19 the instances where it has




Figure 3.6: Some composite gallery possibilities for I_1

3 or 4, and Figure 3.20 the instances where it has 5 or 6. Figure 3.21 is the combined results for n chambers after the departure edge, for $n \geq 0$.

Case 2: $w = f$. The combined results for this case are in Figure 3.22.

Again, if w is some other element of W , then the results are rotations of one of the above cases by 120° and 240° about the center of C_M . Conglomerating all results from I_1 and I_2 gives the results pictured in Figure 3.23. The chambers which are shaded more darkly in this figure are the chambers $w^{-1}bwC_M$ for $w \in W$.

We will not compute other infinite classes of type-edge pairs here, but we will describe how the collection of type-edge pairs can be divided into infinite classes. For each corridor c_i , we get two infinite classes, $I_{c_i}^1$ and $I_{c_i}^2$. The SMG types in each of these classes are those corresponding to SMGs whose terminus is in c_i . If η_1 and η_2 are the two possible angles of edges of departure of these SMGs, then $I_{c_i}^j$ has type-edge pairs whose edge of departure component has angle η_j . Note that I_2 is the union of two infinite classes of this kind, namely $I_{c_4}^1$ and $I_{c_4}^2$.

If G is an SMG type corresponding to an SMG whose final chamber is in one of the regions R_i or r_i , then G has a turning point. The edge of departure for G could be before, after, or at the turning point. However, we need not consider type-edge pairs for which the edge of departure is after or at the turning point, because Γ_E^1 for these type-edge pairs is the same as that arising from some type-edge pair in some $I_{c_i}^j$.

So for each i we have $I_{R_i}^1$ and $I_{R_i}^2$ with SMG types corresponding to an SMG whose final chamber is in R_i . The allowed edges of departure are those before the




Figure 3.7: Class I_1 , where Γ_E^1 has 3 chambers after the turning edge




Figure 3.8: Class I_1 , where Γ_E^1 has 5 chambers after the turning edge




Figure 3.9: Class I_1 , where Γ_E^1 has 7 chambers after the turning edge





Figure 3.10: Summary of results for I_1 , an odd number of chambers after the turning edge

Figure 3.11: More composite gallery possibilities for I_1

turning edge, with angle specified by the superscript. Note that I_1 is of this type. We define $I_{r_i}^1$ and $I_{r_i}^2$ analogously for $i = 1, 2, 3$.

When one conglomerates the results from all infinite classes of composite galleries for

$$b = \begin{pmatrix} \pi^3 & 0 & 0 \\ 0 & \pi^{-1} & 0 \\ 0 & 0 & \pi^{-2} \end{pmatrix},$$

one gets an upper bound set S_1 . It turns out that infinite classes other than those subsumed by I_1 and I_2 do not contribute any chambers beyond those contributed by I_1 and I_2 . So Figure 3.23 is the complete superset for b with $\alpha = 3$ and $\beta = -1$.

When one conglomerates the results of the classes I_1 and I_2 for

$$b = \begin{pmatrix} \pi^2 & 0 & 0 \\ 0 & \pi^0 & 0 \\ 0 & 0 & \pi^{-2} \end{pmatrix},$$

one gets the chambers pictured in Figure 3.24. The results of other infinite classes have not been computed in this case, but it is reasonable to expect that they would not contribute any chambers additional to those contributed by I_1 and I_2 . This will be discussed further later.




Figure 3.12: Class I_1 , where Γ_{E_1} has 2 chambers after the turning edge




Figure 3.13: Class I_1 , where Γ_{E_1} has 4 chambers after the turning edge




Figure 3.14: Class I_1 , where Γ_{E_1} has 6 chambers after the turning edge







Figure 3.15: Summary of results for I_1 , an even number of chambers after the turning edge

Figure 3.16: Some composite gallery possibilities for I_2 Figure 3.17: More composite gallery possibilities for I_2 Figure 3.18: Class I_2 , where Γ_{E_1} has 0, 1 or 2 chambers after the departure edge, $w = 1$




Figure 3.19: Class I_2 , where Γ_{E_1} has 3 or 4 chambers after the departure edge, $w = 1$




Figure 3.20: Class I_2 , where Γ_{E_1} has 5 or 6 chambers after the departure edge, $w = 1$




Figure 3.21: Summary of results for I_2 , $w = 1$




Figure 3.22: Summary of results for I_2 , $w = f$




Figure 3.23: Summary of results for I_1 and I_2

When one conglomerates the results of the classes I_1 and I_2 for

$$b = \begin{pmatrix} \pi^2 & 0 & 0 \\ 0 & \pi^1 & 0 \\ 0 & 0 & \pi^{-3} \end{pmatrix},$$

one gets the chambers pictured in Figure 3.25. Again, other infinite classes have not been computed, but it is reasonable not to, as will be discussed later.

When one conglomerates the results of the classes I_1 and I_2 for

$$b = \begin{pmatrix} \pi^4 & 0 & 0 \\ 0 & \pi^0 & 0 \\ 0 & 0 & \pi^{-4} \end{pmatrix},$$

one gets the chambers pictured in Figure 3.26. Again, other infinite classes have not been computed, but it is reasonable not to.

When one conglomerates the results of the classes I_1 and I_2 for $b = 1$, one gets the chambers pictured in Figure 3.27. We have also computed the results of all other infinite classes of composite galleries in the $b = 1$ case, as we did for the $\alpha = 3$, $\beta = -1$ case. Again, one can see (after the fact) that the infinite classes beyond those subsumed by I_1 and I_2 do not contribute any additional chambers.

The computation of the results of I_1 and I_2 for $b = 1$ is a little different from the cases done thus far, all of which are similar to the example $\alpha = 3$, $\beta = -1$ that was done in detail. The general shapes for composite galleries for $b = 1$ with $(t, e) \in I_1$ are shown in Figure 3.28 for $w = f$ and Figure 3.29 for $w = 1$. The general shapes for $b = 1$ with $(t, e) \in I_2$ are shown in Figure 3.30 for $w = 1$. The $w = f$ for I_2 galleries would be similar. All these galleries are different in general shape from their $\alpha = 3$, $\beta = -1$ counterparts. Note that the I_2 galleries are minimal, and therefore would be easy to fold.

When one conglomerates the results of the classes I_1 and I_2 for

$$b = \begin{pmatrix} \pi^4 & 0 & 0 \\ 0 & \pi^{-2} & 0 \\ 0 & 0 & \pi^{-2} \end{pmatrix},$$

one gets the chambers pictured in Figure 3.31.

The computation of the results of I_1 and I_2 for this last value of b is also a little different from the cases done thus far because $b \neq 1$, but $\alpha + 2\beta = 0$. This is to say that bC_M sits along the bottom edge of the positive Weyl chamber. We will call any b with $\alpha + 2\beta = 0$ or $\alpha + 2\beta = 2\alpha + \beta$ *degenerate*. The values of bC_M for degenerate b are pictured in Figure 3.32. Note that the condition $\alpha + 2\beta = 2\alpha + \beta$ corresponds to the bC_M along the top-left boundary of the positive Weyl chamber.




Figure 3.24: Result for $\alpha = 2, \beta = 0$




Figure 3.25: Result for $\alpha = 2, \beta = 1$




Figure 3.26: Result for $\alpha = 4$, $\beta = 0$




Figure 3.27: Result for $b = 1$




Figure 3.28: General shape of composite galleries for $b = 1$, $w = f$, I_1




Figure 3.29: General shape of composite galleries for $b = 1$, $w = 1$, I_1




Figure 3.30: General shapes of composite galleries for $b = 1$, $w = 1$, I_2





Figure 3.31: Result for $\alpha = 4$, $\beta = -2$

Figure 3.32: bC_M for degenerate b

The general shapes of composite galleries for $b \neq 1$, $\alpha + 2\beta = 0$, $(t, e) \in I_1$ are pictured in Figure 3.33 for $w = f$ and Figure 3.34 for $w = 1$. The composite galleries for I_2 in these cases are similar enough in general shape to those of non-degenerate b that folding considerations are essentially the same.

In the case that $\alpha + 2\beta = 2\alpha + \beta$, $b \neq 1$, the composite galleries for I_2 become different in general shape from those of non-degenerate b , but those for I_1 are similar.

Even though the general shape of composite galleries is different in each of the four categories: b non-degenerate, $b = 1$, $\alpha + 2\beta = 0$ but $b \neq 1$, $\alpha + 2\beta = 2\alpha + \beta$ but $b \neq 1$, there is an efficient way of computing the folding results of these galleries that turns out to be very similar in all four cases. This efficient method is somewhat different from that used previously to compute the $\alpha = 3$, $\beta = -1$ example (although the efficient method will also apply to that example), and will be described in Section 3.1.3.

The lines superimposed on the patterns in Figures 3.23, 3.24, 3.25, 3.26, 3.27, and 3.31 are only there to make the patterns easier to view and to compare. The chambers which are shaded more darkly in Figure 3.23 are the chambers $w^{-1}bwC_M$ for $w \in W$. These are not shaded on the other figures because one can tell where they are by analogy (or by an easy computation). Note that for Figure 3.27 and Figure 3.31, some of the $w^{-1}bwC_M$ are equal to each other. These are degenerate cases, and will be discussed further in Section 3.1.3.




Figure 3.33: General shape of composite galleries for $b \neq 1$ with $\alpha + 2\beta = 0$, $w = f$, I_1




Figure 3.34: General shape of composite galleries for $b \neq 1$ with $\alpha + 2\beta = 0$, $w = 1$, I_1

It is reasonable to expect that other infinite classes would not contribute further to the results in Figures 3.24, 3.25, or 3.26, and 3.31 as they do not contribute further to the results in Figures 3.23 and 3.27.

It is easy to generalize what the results of classes I_1 and I_2 would be for any of the b we are working with (listed in the beginning of Section 3.1). This will also be done in detail in Section 3.1.3. Since the computations involved in discovering the total result of all infinite classes of type-edge pairs are very lengthy for any given b , no computation general to all b has been done. It seems very reasonable that in any case the total results would be the same as the I_1 and I_2 results together, as mentioned previously.

3.1.3 Relationship Between $X_w(1\sigma)$ and $X_w(b\sigma)$, and an Efficient Way of Computing Supersets

In this section we try to illuminate the cause of some of the similarities among the supersets S_1 for different values of b . We do this by discussing a more efficient method of computing the infinite classes I_1 and I_2 that were computed in Section 3.1.2. The basic idea is as follows. Let Γ_{E_1} and Γ_{E_2} be two composite galleries (constructed in Section 3.1.1) that need to be folded in order to compute one of the supersets of Section 3.1.2. Let p_i for $i = 1, 2$ be the number of chambers appearing in $\Gamma_{E_i}^1$ between the edge of departure and the turning edge, let q_i for $i = 1, 2$ be the number of chambers appearing in $\Gamma_{E_i}^1$ after the turning edge, and let w_i for $i = 1, 2$ be elements of W such that $a_i w_i C_M = \rho(E_i)$ for some $a_i \in T(F)$. If $p_1 = p_2$ and $w_1 = w_2$, then the folding results of Γ_{E_1} and Γ_{E_2} can be computed in very similar ways. In fact, for most fixed values of $p_1 = p_2$, for any fixed values of $w_1 = w_2$, and for most b , one can compute folding results for arbitrary q_i all at once. This is done by replacing all the composite galleries with the given p and w with a single “half-infinite gallery” which one can show in advance will have the same folding results as all these composite galleries together. The folding results of the “half-infinite gallery” are easy to compute.

At the end of Section 3.1.1 we defined the *transition type* between two chambers that share an edge, and the *type* of a non-stuttering gallery G in $\mathcal{B}_\infty$. We also need the following definitions.

Definition 3.1.3. A *finite gallery type* is, equivalently,

- 1) An initial chamber $C_0 \subseteq A_M$ and a finite sequence of transition types, which we will call $\{t_1, t_2, \dots, n\}$
- 2) A non-stuttering finite gallery in A_M .

Definition 3.1.4. A *left-infinite gallery type* is, equivalently,

- 1) A terminal chamber $C_0 \subseteq A_M$ and a left-infinite sequence $\{\dots, t_{-3}, t_{-2}, t_{-1}, t_0\}$ of transition types.
- 2) A left-infinite non-stuttering gallery in A_M .

Let $\{C_M; t_1, t_2, \dots, t_n\}$ be a finite gallery type whose initial gallery is C_M . Let $C_M = C_0, C_1, \dots, C_n$ be the corresponding non-stuttering gallery in A_M . We call this the *standard folding* of the gallery type $\{C_M; t_1, t_2, \dots, t_n\}$. There are *non-standard foldings* of $\{C_M; t_1, t_2, \dots, t_n\}$ determined as follows. We first assume inductively that C_i is determined. Note that the base case, $C_0 = C_M$ actually is determined. Then let L_{i+1} be the wall containing the edge of C_i that does not contain the vertex of type t_{i+1} . If C_M and C_i are on the same side of L_{i+1} , then let C_{i+1} be the reflection of C_i about L_{i+1} . If C_M and C_i are on opposite sides of L_{i+1} , then let C_{i+1} be either C_i itself, or the reflection of C_i about L_{i+1} . The choices that arise in this process may lead to many different non-standard foldings of $\{C_M; t_1, t_2, \dots, t_n\}$.

Definition 3.1.5. The *folding results* of $\{C_M; t_1, t_2, \dots, t_n\}$ are the collection of possible final chambers C_n that can arise from all possible (standard and non-standard) foldings of $\{C_M; t_1, \dots, t_n\}$.

We now let $\{\dots, t_{-3}, t_{-2}, t_{-1}, t_0; C_0\}$ be a left-infinite gallery type, and we let $\{\dots, C_{-3}, C_{-2}, C_{-1}, C_0\}$ be the corresponding non-stuttering gallery in A_M . Then it is easy to see that for any n , $\{C_{-n}; t_{-n+1}, t_{-n+2}, \dots, t_0\}$ is a finite gallery type. Let R_n be the folding results of this finite gallery type. Let $R = \cup_{n=1}^{\infty} R_n$. We call R the *folding results* of $\{\dots, t_{-3}, t_{-2}, t_{-1}, t_0; C_0\}$.

Each type-edge pair $(t, e) \in I_1 \cup I_2$ gives Γ_x a composite gallery, and the folding results of that type-edge pair, $S_{(t,e)}$ are the folding results of the finite gallery type that Γ_x represents. In fact, we frequently have $S_{(t_1, e_1)} = S_{(t_2, e_2)}$ for $(t_1, e_1) \neq (t_2, e_2)$. If $(t, e) \in I_1$, then there are three characteristics of (t, e) that are clearly relevant to the type of $x^{-1}\Gamma_x$, and therefore relevant to $S_{(t,e)}$. First, the number p of chambers between e and the turning edge of t will have significance. Second, the number q of chambers after the turning edge of t will have significance. And third, the Weyl group element $w \in W$ such that $awC_M = C_n$ will have significance, where a is a diagonal matrix, and C_n is the last chamber of t . It is easy to see that if $p_{(t_1, e_1)} = p_{(t_2, e_2)}$, $q_{(t_1, e_1)} = q_{(t_2, e_2)}$, and $w_{(t_1, e_1)} = w_{(t_2, e_2)}$ then $S_{(t_1, e_1)} = S_{(t_2, e_2)}$, for $(t_i, e_i) \in I_1$. Let $R_{p,q,w}^1$ be this set.

For $(t, e) \in I_2$, we only have q , the number of chambers after e in t ; and w . If $q_{(t_1, e_1)} = q_{(t_2, e_2)}$, and $w_{(t_1, e_1)} = w_{(t_2, e_2)}$ for $(t_i, e_i) \in I_2$, then $S_{(t_1, e_1)} = S_{(t_2, e_2)}$. Let $R_{q,w}^2$ be this set.

For fixed p, w , let $R_{p,w}^1 = \cup_q R_{p,q,w}^1$, and for fixed w let $R_w^2 = \cup_q R_{q,w}^2$. We will show that $R_{p,w}^1$ and R_w^2 can usually be computed as the folding results of some half-infinite gallery type. Let $r \in W$ be rotation by 120° counterclockwise, and let $f \in W$ be the reflection about the horizontal line through v_M . For b with $\alpha + 2\beta > 0$, and given $p \geq 1$ odd, we define $\Omega_1^2, \Omega_f^2, \Omega_{p,1}^1$ and $\Omega_{p,f}^1$ to be the half-infinite galleries shown in Figure 3.35 (This figure shows the specified half-infinite galleries for a particular choice of b , namely $\alpha = 4$ and $\beta = -1$. But it is clear how the definition would

work for other b with $\alpha + 2\beta > 0$.) The half-infinite galleries Ω_r^2 , Ω_{rf}^2 , $\Omega_{p,r}^1$ and $\Omega_{p,rf}^1$ are rotations of those in Figure 3.35 by 240° counterclockwise about the center point of C_M , and $\Omega_{r^2}^2$, $\Omega_{r^2f}^2$, Ω_{p,r^2}^1 and Ω_{p,r^2f}^1 are rotations of those in Figure 3.35 by 120° counterclockwise about the center point of C_M . If $\alpha + 2\beta = 0$ then $\Omega_1^2 = \Omega_f^2$, $\Omega_r^2 = \Omega_{rf}^2$, and $\Omega_{r^2}^2 = \Omega_{r^2f}^2$ (the picture would be a degenerate version of Figure 3.35).

Proposition 3.1.2. If $\alpha + 2\beta > 0$ or $b = 1$, then $R_{p,w}^1$ is the folding results of $\Omega_{p,w}^1$, and R_w^2 is the folding results of Ω_w^2 . If $\alpha + 2\beta = 0$ and $b \neq 1$, then $(\cup_p R_{p,w}^1) \cup R_w^2$ is the union of the folding results of the $\Omega_{p,w}^1$ for all p , and the folding results of Ω_w^2 . It is still true that R_w^2 is the folding results of Ω_w^2 .

Before we prove this proposition, we establish some more terminology. We first let $\{C_M; t_1, t_2, \dots, t_n\}$ be a finite gallery type. Let $C_0 = C_M, C_1, \dots, C_m$ be a (standard or non-standard) folding of the sub-gallery type $\{C_M; t_1, t_2, \dots, t_m\}$, where $m < n$. Consider the finite gallery type $\{C_m; t_{m+1}, \dots, t_n\}$, and let $C_m = D_m$, D_{m+1} , $D_{m+2}, \dots, D_n$ be the standard folding. If D_{m+j} , for $1 \leq j \leq n - m$ is on the same side of the edge between D_{m+j} and D_{m+j-1} as C_M , then we say $m + j$ is a *choice point given the history* $C_M, C_1, \dots, C_m$. Note that this all applies even if $m = 0$. Also note that if $1 \leq k \leq n$, then k could be a choice point given some histories, but not given others. We now prove the proposition.

Proof. It suffices to consider $w = 1$ and $w = f$. In fact, we will only consider $w = 1$, since the $w = f$ case is similar. We first assume that $\alpha + 2\beta > 0$ or $b = 1$, and that $(t, e) \in I_1$. Suppose $x \in SL_3(L)$ has SMG with type t and edge of departure e . Also suppose (t, e) has characteristics p, q , and $w = 1$. Let Γ_x be the composite gallery. So in the notation of Section 3.1.1, $\Gamma_x = \Gamma_E^1 \cup \Gamma_E^2 \cup \Gamma_E^3$, where $E = xC_M$. Consider the choice points given history $C_M = C_0$. These all lie in Γ_E^3 , because of the shape of Γ_x (illustrated for $\alpha = 3, \beta = -1, q = 11$ and $p = 7$ in Figure 3.3). The choice points given history $C_M = C_0$ for the same b and x that were used in Figure 3.3 are illustrated in Figure 3.36. Note that $\Omega_{p,1}^1$ ends at the same chamber that the standard folding of Γ_x ends at, and note that the choice points in Γ_x that are after or at the turning edge of Γ_E^3 are also choice points in $\Omega_{p,1}^1$. This is illustrated for the example $\alpha = 3, \beta = -1, w = 1, p = 7$ and $q = 11$ in Figure 3.37. Note also that if i is a choice point in Γ_E^3 that occurs before the turning edge of Γ_E^3 , and if L_i is the wall containing the edge corresponding to i , then L_i also passes through $\Omega_{p,1}^1$. Reflecting the section of Γ_E^3 that occurs after i about L_i gives rise to a finite gallery type with no choice edges additional to those obtained by reflecting the portion of $\Omega_{p,1}^1$ that occurs after L_i about L_i . This is illustrated in Figure 3.38. Therefore, the folding results of $\Omega_{p,1}^1$ contain $R_{p,1}^1$.




Figure 3.35: Ω^2_1 , Ω^2_f , $\Omega^1_{p,1}$ and $\Omega^1_{p,f}$

The figure shows a triangular lattice. A shaded region, labeled C_M , is located in the lower-left. A boundary line, labeled $\Omega_{7,1}^1$, is shown in the upper-right. The boundary line is a zigzag line that follows the edges of the triangles. The shaded region C_M is a large, irregular shape that is shaded gray. The boundary line $\Omega_{7,1}^1$ is a black line that follows the edges of the triangles.

Figure 3.37: Choice points for $\Omega_{p,1}^1$ and Γ_x




Figure 3.38: How to address choice points of Γ_x that occur before the turning point of Γ_E^3

We must now show that $R_{p,1}^1$ contains the folding results of $\Omega_{p,1}^1$. It suffices to show that every choice point in $\Omega_{p,1}^1$ is also a choice point in some Γ_x for some xC_M with type-edge pair having characteristics p , $w = 1$, and some q . But this can be accomplished by letting q get arbitrarily large.

To see that R_1^2 is the folding result of Ω_1^2 for $\alpha + 2\beta \neq 2\alpha + \beta$ or for $b = 1$, proceed as follows. First note that Γ_x for x with $p = 0$ is minimal, and the final gallery of $\rho(x^{-1}\Gamma_x)$ is bC_M . This is also the final gallery of the standard folding of Ω_1^2 . Then let the chambers in the standard folding of Ω_1^2 be called $\{\dots, D_3, D_2, D_1\}$, where $D_1 = bC_M$. Let the wall between D_i and D_{i+1} be called L_i . Let G_i be the left-infinite gallery in A_M that one gets by reflecting the chambers D_j for $j \leq i$ across L_i . Note that if $p \geq 1$, the only choice points in Γ_x are duplicated in G_p . This shows that R_1^2 is equal to the set of folding results of Ω_1^2 .

The $\alpha + 2\beta = 0$ for $b \neq 1$, $(t, e) \in I_1$ case is more complicated, but makes use of the same ideas, and is therefore omitted. The $2\alpha + \beta = \alpha + 2\beta$ case for $b \neq 1$, $(t, e) \in I_2$ is omitted for the same reason. $\square$

This gives a more efficient way to compute the folding results of all the type-edge pairs in $I_1 \cup I_2$ for any fixed b , and also illustrates, to some extent, the source of the similarity between the sets S_1 for different values of b .

3.1.4 A Method Suggested by Rapoport and Kottwitz

The set S_1 of chambers in A_M is, *a priori*, only a superset of the solution set S that we are seeking. In this section we give the results of a methodology suggested by Rapoport and Kottwitz for producing a subset $S_2 \subseteq S$. In other words, this method will show that certain chambers in A_M must be contained in S .

The method in this section is only effective in the case $b = 1$. In this case, techniques that will be described in the next section will serve to enlarge S_2 to the point that it becomes equal to S_1 , and therefore to S . In the $b \neq 1$ cases, we will use other methods to arrive at a subset S_2 (see Section 3.1.6).

Let $a \in SL_3(F)$ be of the form

$$a = \begin{pmatrix} \pi^m & 0 & 0 \\ 0 & \pi^n & 0 \\ 0 & 0 & \pi^{-m-n} \end{pmatrix},$$

where there are no conditions on m and n . Let w be one of the following matrices:

$$r = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad ; \quad r^2 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}.$$

Therefore, w represents a 3-cycle in W , the finite Weyl group. According to a result of Kottwitz, the matrix aw belongs to a *basic* σ -conjugacy class if there is some l such that $aw\sigma(aw)\sigma^2(aw)\dots\sigma^{l-1}(aw)$ is central in $SL_3(L)$ [6], [7]. See [6] and [7] for a definition of *basic*, but for the present considerations, one only needs to know that the σ -conjugacy class containing $b = 1$ is the only basic σ -conjugacy class of $SL_3(L)$.

Lemma 3.1.3. There exists an l such that $aw\sigma(aw)\sigma^2(aw)\dots\sigma^{l-1}(aw)$ is central for any choice of m, n .

Proof. First we note that $\sigma(aw) = aw$, since $aw \in SL_3(F)$. So we want to show that $(aw)^l$ is central for some l . But

$$\begin{aligned} (aw)^l &= awaw(aw)^{l-2} \\ &= awaw^{-1}w^2(aw)^{l-2} \\ &= aa^w w^2(aw)^{l-2}, \end{aligned}$$

where the superscript w denotes w acting on a by conjugation. Proceeding, we have

$$\begin{aligned} aa^w w^2(aw)^{l-2} &= aa^w w^2 aw(aw)^{l-3} \\ &= aa^w w^2 aw^{-2} w^3(aw)^{l-3} \\ &= aa^w a^{w^2} w^3(aw)^{l-3} \\ &= \dots \\ &= aa^w a^{w^2} a^{w^3} \dots a^{w^{l-1}} w^l. \end{aligned}$$

Choosing $l = 3$, we get $aa^w a^{w^2} w^3 = 1$, which is central.

Since w is a 3-cycle, conjugating by w serves to permute the diagonal entries of a in a cyclic way. This is why $aa^w a^{w^2} w^3$ is central. $\square$




Figure 3.39: Main results of the method suggested by Rapoport and Kottwitz

So the matrices aw are in the σ -conjugacy class of 1 for any choice of m, n . Therefore, the double- I -cosets $\{IawI : m, n \in \mathbb{Z}\}$ all meet the identity σ -conjugacy class non-trivially. These double- I -cosets are pictured in Figure 3.39.

With certain conditions on m and n , it is also possible for $(aw)^l$ to be central for some l if w is not a 3-cycle in W :

Lemma 3.1.4. If $a = 1$ then there exists l such that $(aw)^l$ is central for any $w \in W$.

Proof. Take $l = 6$. □

The implication of this is that the double- I -cosets corresponding to the chambers in Figure 3.40 all intersect the σ -conjugacy class of 1. Let f be as in Section 3.1.2.

Lemma 3.1.5. If $m = 0$ then $aa^w = 1$ for $w = f$. If $m = -n$ then $aa^w = 1$ for $w = rf$. If $n = 0$ then $aa^w = 1$ for $w = r^2f$.

Proof. Just compute the relevant matrix products. □

The implication of this lemma is that the double- I -cosets corresponding to the chambers in Figure 3.41 all intersect the σ -conjugacy class of 1. Note, in addition, that we have a concrete element of the intersection of each of these double- I -cosets with $\{x^{-1}\sigma(x) : x \in SL_3(L)\}$.




Figure 3.40: The $a = 1$ results of the method suggested by Rapoport and Kottwitz





Figure 3.41: Results from the method suggested by Rapoport and Kottwitz under the circumstances of Lemma 3.1.5

Figure 3.42: The choice D_1 of xC_M

It is easy to see that $(aw)^l$ is not central for any l , unless we are in one of the cases enumerated above.

It is clear how one would go about formulating and proving the lemmas of this section for SL_n . One would get results for every possible cycle decomposition in $W = S_n$, and the “most” result chambers ($\mathbb{Z}^{n-1}$ worth of them) would arise from the n -cycles, since then a is allowed to be arbitrary. At the other end of the spectrum, the finitely many Weyl group images of the main chamber arise from $a = 1$. In subsequent sections, the results from this section will also be generalized to Sp_{2n} and G_2 .

3.1.5 A Subset of the Solution Set for $b = 1$

Let S_2 be the collection of all double- I -cosets shown in the previous section to meet $\{x^{-1}\sigma(x) : x \in SL_3(L)\}$. So S_2 is the union of the chambers pictured in Figures 3.39, 3.40, and 3.41. We know that $S_2 \subseteq S \subseteq S_1$, and in this section we enlarge S_2 in such a way that it remains a subset of S . The enlarged S_2 will turn out to be equal to S_1 . Note that this section concerns itself only with $b = 1$.

Let $w \in SL_3(L)$ be one of the matrices r, r^2 from the previous section. Let a be as in the previous section. Then we know $\tilde{b} = aw$ is σ -conjugate to $b = 1$, so $\{x^{-1}\sigma(x) : x \in SL_3(L)\} = \{x^{-1}\tilde{b}\sigma(x) : x \in SL_3(L)\}$, and also $\{Ix^{-1}\sigma(x)I : x \in SL_3(L)\} = \{Ix^{-1}\tilde{b}\sigma(x)I : x \in SL_3(L)\}$. But we know that the chamber in A_M corresponding to $Ix^{-1}\tilde{b}\sigma(x)I$ is $\rho(x^{-1}\tilde{b}\sigma(x)C_M)$. Just as in Section 3.1.1, let Γ_x^1 be the part of the SMG from C_M to xC_M that is not in A_M , let $\Gamma_x^3 = \tilde{b}\sigma(\Gamma_x^1)$, and, if e is the only edge of Γ_x^1 in A_M , let Γ_x^2 be a minimal gallery from e to $\tilde{b}\sigma(e) = \tilde{b}e$. If $\Gamma_x = \Gamma_x^1 \cup \Gamma_x^2 \cup \Gamma_x^3$, then the possible foldings of galleries of the same type as $x^{-1}\Gamma_x$ give possible candidates for additions to the set S_2 . For some choices of xC_M there is only one possible folding of galleries of the same type as $x^{-1}\Gamma_x$, and the final chamber of this folding is not already in S_2 . Such a situation would give an additional chamber to add to S_2 .

For instance, we can choose xC_M to be the chamber in Figure 3.42 that is adjacent to C_M , but not in A_M . If $w = r^2$ and




Figure 3.43: The resulting Γ_x for D_1 and a particular choice of aw




Figure 3.44: The result of folding galleries of the same type as $x^{-1}\Gamma_x$ for the Γ_x pictured in Figure 3.43

$$a = \begin{pmatrix} \pi^3 & 0 & 0 \\ 0 & \pi^{-1} & 0 \\ 0 & 0 & \pi^{-2} \end{pmatrix},$$

then Γ_x is shown in Figure 3.43. Any gallery of the same type as $x^{-1}\Gamma_x$ folds down into A_M in the same way, so we know that $Ix^{-1}\tilde{b}\sigma(x)I$ corresponds to the chamber shown in Figure 3.44, which was not obtained by the methods of the previous section.

We now keep w as above, but let a take the general form

$$a = \begin{pmatrix} \pi^m & 0 & 0 \\ 0 & \pi^n & 0 \\ 0 & 0 & \pi^{-m-n} \end{pmatrix}.$$

For these $\tilde{b} = aw$ we can compute the possible ways that galleries of the same type as $x^{-1}\Gamma_x$ can fold into A_M , where, as before, x is chosen so that xC_M is the chamber D_1 pictured in Figure 3.42. One can easily see that for each m, n , there is only one way that such galleries can fold down. The chambers obtained through this process are marked in Figure 3.45. Many of these did not arise from the methods of the previous section.

We could also choose x such that xC_M is one of the chambers D_2, D_3 in Fig-





Figure 3.45: Results for $xC_M = D_1$ and $w = r^2$ or for $xC_M = D_1$ and $w = r$

Figure 3.46: The choices D_2 and D_3 of xC_M

w	xC_M	Figure containing results
r^2	D_1	Figure 3.45
r^2	D_2	Figure 3.47
r^2	D_3	Figure 3.48
r	D_1	Figure 3.45
r	D_2	Figure 3.47
r	D_3	Figure 3.48

Table 3.1: Which figure contains which results

ure 3.46, and we could choose $w = r$. For each of these possible choices, we could make considerations similar to the above. The results of this process are pictured in subsequent figures according to Table 3.1.

If we combine all these results with those of the previous section, we get the chambers pictured in Figure 3.49. Note that this combined set of chambers is still not equal to S_1 .

If $\tilde{D}_4$, $\tilde{D}_5$, and $\tilde{D}_6$ are the chambers in A_M labeled in Figure 3.50, and if $g_4, g_5, g_6 \in I$ are chosen such that $g_i E_i \neq E_i$, then we can choose x such that xC_M is any of the $D_i = g_i \tilde{D}_i$. For each of these xC_M , we could make considerations similar to the above for $w = r, r^2$. The results of this process include all the chambers in Figure 3.27 that are not present in Figure 3.49. This proves that the chambers in Figure 3.27 represent the exact collection of double- I -cosets that intersect non-trivially with the σ -conjugacy class of 1.

Rather than using xC_M a distance of two chambers from C_M , it is also possible to obtain the chambers in Figure 3.27 that are not in Figure 3.49 in the following way. Let $\tilde{b}$ now be such that $\tilde{b}C_M$ is one of the chambers in Figure 3.49 that was not obtained using the methods of the previous section. Let $xC_M \notin A_M$ be D_1 , D_2 or D_3 , and compute as previously. This gives rise to the chambers in Figure 3.27 that are not in Figure 3.49. Note that this constitutes an iteration of our “length




Figure 3.47: Results for $xC_M = D_2$ and $w = r^2$ or for $xC_M = D_2$ and $w = r$




Figure 3.48: Results for $xC_M = D_3$ and $w = r^2$ or for $xC_M = D_3$ and $w = r$




Figure 3.49: Combination of all results referred to in Table 3.1




Figure 3.50: Some choices for $x C_M$ that are two chambers away from C_M




Figure 3.51: A Useful Lemma

one appendage” methods, and gives rise to the same additional chambers that our “length two appendage” methods did.

The methods in this section assume the existence of a large and well-distributed set S_2 . If one has a smaller S_2 , the enlargement effort will be less effective. One could apply the methods of this section to σ -conjugacy classes other than $b = 1$, but one would have to somehow come up with a reasonably large S_2 first.

3.1.6 A Geometric Construction of a Subset of the Solution Set

In this section, we prove using geometric methods that the chambers in the superset S_1 produced in Section 3.1.2 that arise from the infinite classes I_1 and I_2 are all actually contained in the solution set S . Since the classes I_1 and I_2 seemed to give rise to all chambers in S_1 , this proves that $S_1 = S$.

We begin with the following very useful lemma.

Lemma 3.1.6. Let G be one of the galleries pictured in Figure 3.51. Let $\tilde{G} = gG$, where $g \in GL_3(L)$ is arbitrary. There exists a unique vertex $\tilde{v}$ such that $\tilde{v}$ is adjacent to each of gv_5 , gv_1 , and gv_M . Here, v_M is the main vertex in the building.

Proof. It suffices to work with $g = 1$. The vertices adjacent to v_M correspond to non-trivial proper subspaces of $(\overline{\mathbb{F}}_q)^3$, where $\mathbb{F}_q$ is the residue field of F (so $\overline{\mathbb{F}}_q$ is the residue field of L). For such a vertex v , let v also denote the corresponding subspace of $(\overline{\mathbb{F}}_q)^3$. If v and w are two vertices adjacent to v_M , then v and w are adjacent to each other if and only if $v \subseteq w$ or $w \subseteq v$ as subspaces. We therefore have the following cases:

Case 1: $\dim(v_1) = 1$. Then $\dim(v_2) = 2$, $\dim(v_3) = 1$, $\dim(v_4) = 2$, $\dim(v_5) = 1$, and we can and must choose $\tilde{v} = \langle v_1, v_5 \rangle$.

Case 2: $\dim(v_1) = 2$. Then $\dim(v_2) = 1$, $\dim(v_3) = 2$, $\dim(v_4) = 1$, $\dim(v_5) = 2$, and we can and must choose $\tilde{v} = v_1 \cap v_5$. $\square$

The idea of this lemma is that whenever we see an arrangement of chambers shaped like one of those in Figure 3.51, we can fill it in uniquely with two more chambers to create a hexagon.

We also need the following definition:

Definition 3.1.6. Given a minimal gallery G , the *parallelogram*, $P(G)$ associated with G is $P(G) = \cap A$, where the intersection is over all apartments A containing G .

Theorem 3.1.7. $P(G) = \cup \tilde{G}$, where the union is over all minimal galleries $\tilde{G}$ that stretch from the first chamber of G to the last chamber of G .

Proof. See [8]. □



Note that $P(G)$ is not necessarily actually a parallelogram. It is a parallelogram with possibly a single chamber removed from either or both of the acute angle corners.



We now discuss the the program that we will follow in the rest of this section. We will begin by restricting our attention to the collection of σ -conjugacy classes consisting of those b with $\alpha + 2\beta \neq 0$, and $b = 1$. We also restrict our attention to type-edge pairs in I_1 . If $x \in SL_3(L)$ gives rise to a type-edge pair $(t, e) \in I_1$, and if b is restricted as specified, then we will define two invariants γ_1 and γ_2 of x . It will turn out that γ_1 and γ_2 determine $\rho(x^{-1}\Gamma_x)$. We will also show that given any γ_1 and γ_2 , we can choose x such that x and b give rise to (t, e) , γ_1 , and γ_2 . This proves that for $b = 1$ or b with $\alpha + 2\beta \neq 0$, all the results in S_1 which come from I_1 are in fact in S . We then use similar methods, with a few added complications, to address $b \neq 1$ with $\alpha + 2\beta = 0$, still focusing our attention on $(t, e) \in I_1$. We then turn to $(t, e) \in I_2$, and we consider separately the cases $b = 1$ or $\alpha + 2\beta \neq 2\alpha + \beta$, and $b \neq 1$ with $\alpha + 2\beta = 2\alpha + \beta$.

As we just mentioned, we start by assuming $(t, e) \in I_1$ and either $b = 1$ or $\alpha + 2\beta \neq 0$. Recall that the associated composite gallery Γ_x is $\Gamma_E^1 \cup \Gamma_E^2 \cup \Gamma_E^3$, where each Γ_E^i is minimal (if $b = 1$ then $\Gamma_E^2 = \emptyset$). Let $P = P(\Gamma_E^1)$, let $P_b = P(\Gamma_E^2)$, and let $P_\sigma = P(\Gamma_E^3)$. It is worth noting that although we never made a specific choice of Γ_E^2 , $P(\Gamma_E^2)$ is well defined (this relies on the fact that we have either $b = 1$ or b with $\alpha + 2\beta \neq 0$). Figure 3.52 has a picture of P , P_b , and P_σ for

$$b = \begin{pmatrix} \pi^3 & 0 & 0 \\ 0 & \pi^{-1} & 0 \\ 0 & 0 & \pi^{-2} \end{pmatrix},$$

and the example type-edge pair pictured in Figure 3.2. By Lemma 3.1.6, we can choose a vertex w_1 adjacent to u_2 , u_3 , and u_4 . This vertex is not necessarily v_1 , although it may be. By repeated application of Lemma 3.1.6, we can choose the vertices w_2 , w_3 , w_4 and w_5 , labeled on Figure 3.53.

Figure 3.52: An example of P , P_b and P_σ Figure 3.53: w_2 , w_3 , w_4 and w_5


Figure 3.54: Multiple rows of the w_i Figure 3.55: Filling in the rest of $\overline{P}$

For an arbitrary $b \neq 1$ with $\alpha + 2\beta \neq 0$, and for any type-edge pair in I_1 , we can choose vertices analogous to the w_i by repeated application of Lemma 3.1.6. Note that if P_b has more than one row of chambers, then we must choose correspondingly many rows of the w_i , one row at a time, from the bottom up. See Figure 3.54. We can also use Lemma 3.1.6 repeatedly to fill in chambers as labeled in Figure 3.55. We refer to the resulting unique construction consisting of P , P_b , and the chambers added in the two above processes as $\overline{P}$. Note that $\overline{P}$ is the intersection of all apartments containing P and P_b , and is also the union of all minimal galleries from the first chamber of Γ_E^1 to the last chamber in Γ_E^2 .

We now define two invariants which describe the way in which $\overline{P}$ is connected to P_σ .

Definition 3.1.7. Let γ_1 denote the number of edges that $\overline{P}$ and P_σ have in common.



So $\gamma_1 \geq 1$, and in the case pictured in Figure 3.52, $1 \leq \gamma_1 \leq 6$.


Figure 3.56: The definition of the invariant γ_2 , part 1

We define the second invariant γ_2 using Figure 3.56, which shows a situation with $\gamma_1 = 2$. Note that for each $i = 1, 2, 3, 4$, the edges e_i of P_σ and h_i of $\overline{P}$ are different, although they look the same in the figure. We pass to Figure 3.57, which is the same as Figure 3.56, only part of P_σ is not shown for clarity. We now define γ_2 for our example situation. It will be clear from this example how γ_2 is defined in general. If there is a chamber C_1 (pictured in Figure 3.58) having h_1 and d_1 as edges, then it is unique, and not contained in P_σ . In this case we require $\gamma_2 \geq 1$, and we get D_1 and D_2 as pictured (in Figure 3.58) by using Lemma 3.1.6. If there is no C_1 as described then we say $\gamma_2 = 0$. Given C_1 , if there is a chamber C_2 connecting D_2 to d_2 , then it is unique. In this case we require $\gamma_2 \geq 2$, and we get D_3 , D_4 , and D_5 , D_6 by using Lemma 3.1.6 twice. If there is no C_2 as described then we say $\gamma_2 = 1$. Given C_2 , we look for C_3 as labeled. We continue in this manner, with the restriction that γ_2 cannot be bigger than either the number of the h_i or the number of the d_i . If γ_2 is equal to the lesser of these numbers, then we say γ_2 is maximal.

For any parallelogram P , let $T(P)$ be the number of edges along the top of P . Let $S(P)$ be the number of edges along the right side of P . Recall that this could be different from the number of edges along the left side of P . Similarly, $T(P)$ could be different from the number of edges along the bottom of P . This is because the “parallelograms” with which we are working are not true parallelograms in the traditional sense. They are parallelograms that may have one chamber removed from either or both of the acute angle corners. Although note that the upper right acute angle of P (and therefore $\overline{P}$) never has a chamber removed; P_b will always be a parallelogram in the traditional sense of the word; and the lower right acute angle of P_σ never has a chamber removed.

Fixing b and a type-edge pair $(t, e) \in I_1$, let $\gamma_1^{\max} = T(P)$. Choose γ_1 such that

Figure 3.57: The definition of the invariant γ_2 , part 2Figure 3.58: The definition of the invariant γ_2 , part 3

Figure 3.59: $T(P) = S(P) = \infty$

$1 \leq \gamma_1 \leq \gamma_1^{\max}$, and choose γ_2 such that $0 \leq \gamma_2 \leq \min\{S(P), T(\bar{P}) - \gamma_1\} = \gamma_2^{\max}(\gamma_1)$. We will show that one can find x such that xC_M gives the type-edge pair (t, e) , and the invariants γ_1 and γ_2 . We begin by making a simplification.

If b is still fixed; (t_1, e_1) and (t_2, e_2) are two type-edge pairs in I_1 ; P_1 and P_2 are the corresponding parallelograms $P(\Gamma_{E_1}^1)$ and $P(\Gamma_{E_2}^2)$; and if $T(P_1) \leq T(P_2)$ and $S(P_1) \leq S(P_2)$, then $\gamma_1^{\max}(P_1) \leq \gamma_1^{\max}(P_2)$. Given $\gamma_1 \leq \gamma_1^{\max}(P_1)$, we also have $\gamma_2^{\max}(P_1, \gamma_1) \leq \gamma_2^{\max}(P_2, \gamma_1)$. To construct P_1 with invariants $\gamma_1(P_1) \leq \gamma_1^{\max}(P_1)$ and $\gamma_2(P_1) \leq \gamma_2^{\max}(P_1, \gamma_1(P_1))$, it would suffice to construct P_2 with the same invariants. One could then chop off part of P_2 to get the desired P_1 . Therefore it suffices to demonstrate the construction of an “infinite parallelogram” P with $T(P) = S(P) = \infty$ with any fixed (and still finite) invariants γ_1 and γ_2 . The equation $T(P) = S(P) = \infty$ is meant to imply a parallelogram that is half-infinite along the top and along the right side as in Figure 3.59

We now discuss notation to be used in constructions to come. Regions of an apartment of the building will always be denoted using Roman letters with subscript. Single chambers will be labeled using Roman letters with subscript, or just using numbers. At several points we will have to graphically represent structures in the building that cannot be embedded into a Euclidean plane. For instance, we will discuss the main apartment A_M , together with a half apartment $A_1^{\frac{1}{2}}$ coming out of a wall K in A_M . We would picture this example as in Figure 3.60, where R_1 is a region in A_M , R_2 is also in A_M , and $\tilde{R}_2 = A_1^{\frac{1}{2}}$. Thus R_2 and $\tilde{R}_2$ are the same region graphically, but different regions in the construction we are describing. The chamber shaded in Figure 3.60 could be denoted 1 if in R_2 or $\tilde{1}$ if in $\tilde{R}_2$. The tilde will be used generally in this way, with constructions marked with a tilde understood to not be in




Figure 3.60: The meaning of a tilde

$\overline{P}$, P_b , or P_σ , unless specifically included.

We now proceed to demonstrate that one can find x such that xC_M gives (t, e) , γ_1 , and γ_2 . We work with two cases.

Case 1: $b = 1$. Let A_1 be an apartment in $\mathcal{B}_1$, and let K be a wall in A_1 (see Figure 3.61). Let $A_1^{\frac{1}{2}}$ be a half-apartment coming out of K that lies in $\mathcal{B}_2$. If γ_1 is as labeled in Figure 3.61, choose region R_1 in $A_1^{\frac{1}{2}}$. Choose region R_2 from $\mathcal{B}_6 \setminus \mathcal{B}_2$. Let $Q_1 = \sigma(R_1)$ and $Q_2 = \sigma(R_2)$. This determines the regions N_1 and $\tilde{r}_1$ by repeated application of Lemma 3.1.6, and therefore determines $r_1 := \sigma(N_1)$. So $\tilde{r}_1$ is attached to N_1 . We must check that the bottom boundaries of r_1 and $\tilde{r}_1$ have no edges in common to ensure that γ_1 is no bigger than intended. But such commonality would imply that for some edge e along the upper boundary of N_1 , we have $\sigma(e) = e$, i.e., $e \in \mathcal{B}_1$. But let G be a minimal gallery through Q_2 from e to some chamber D in $Q_1 \subseteq \mathcal{B}_2$. The galleries G and $\sigma^2(G)$ are both galleries of the same type from e to D , so therefore $G = \sigma^2(G)$. But G passes through $Q_2 \subseteq \mathcal{B}_6 \setminus \mathcal{B}_2$, so $G \neq \sigma^2(G)$, and we have a contradiction.

We now choose R_3 arbitrarily. This determines $Q_3 = \sigma(R_3)$, and $\tilde{r}_2$, N_2 , $\tilde{C}_2$, $\tilde{E}_2$ by Lemma 3.1.6. The region $\tilde{r}_2$ is attached on its right hand side to $\tilde{r}_1$. Choose $C_2 \neq \tilde{C}_2$ to ensure that γ_2 is no bigger than intended. Fill out the rest of P arbitrarily.

Case 2: $b \neq 1$. We break this case into subcases:

Subcase 1: $\gamma_1 + T(P_b) - 1 \geq \gamma_1 + \gamma_2$, and if equality holds then $\gamma_2 \neq 0$. This situation is pictured in Figure 3.62. The inequality implies that c is at least as far left as a in that figure, and if $a = c$ then $\gamma_2 \neq 0$. Let K be a wall in A_M , and let $A_1^{\frac{1}{2}}$ be a half apartment coming out of K . Then $b\sigma(A_1^{\frac{1}{2}})$ is a half apartment coming out of $b\sigma(K) = bK$. Choose Q_1 and $\tilde{r}_1$ from $b\sigma(A_1^{\frac{1}{2}})$. Let $R_1 = \sigma^{-1}b^{-1}Q_1$. Then

Figure 3.61: Existence of all values of γ_1 and γ_2 , case 1

R_1 is in $A_1^{\frac{1}{2}}$, so $L_1 \subseteq \overline{P}$ is in A_M , and Q_1 and $\tilde{r}_1$ are attached to $\overline{P}$ along $b\sigma(K)$. Choose r_1 the same shape as $\tilde{r}_1$, and attached to Q_1 along the line between Q_1 and $\tilde{r}_1$, but sharing no chambers in common with $\tilde{r}_1$. This ensures that none of the chambers in r_1 is attached to L_1 , and therefore that γ_1 is no bigger than intended. Let $N_1 = \sigma^{-1}b^{-1}(r_1)$. We can fill in L_2 by repeated application of Lemma 3.1.6. Note that L_2 may be empty if $\gamma_2 = 0$. In this case, r_1 , $\tilde{r}_1$ and $\tilde{r}_2$ would also be empty. If $\gamma_2 > 0$, choose $\tilde{r}_2$ attached to L_1 , or attached to L_2 if $c = a$ (and in this case we have $L_2 \neq \emptyset$, since $\gamma_2 \neq 0$). This is done by repeated application of Lemma 3.1.6. Choose Q_2 such that there is no chamber attaching it to $\tilde{r}_2$ (we know this can be done by the uniqueness statement in Lemma 3.1.6). Now let $R_2 = \sigma^{-1}b^{-1}Q_2$. This ensures that γ_2 is no bigger than desired. We now choose the rest of P arbitrarily.

Subcase 2: $\gamma_1 + T(P_b) - 1 = \gamma_1 + \gamma_2$ and $\gamma_2 = 0$. In this case, $T(P_b) = 1$. The situation is pictured in Figure 3.63. Let $A_1^{\frac{1}{2}}$ be a half apartment coming out of K , as before. Choose R_1 in $A_1^{\frac{1}{2}}$, and let $Q_1 = b\sigma(R_1)$. We see that $L_1 \subseteq A_M$. Choose R_2 attached to R_1 . This determines $Q_2 = b\sigma(R_2)$, which in turn determines $\tilde{D}_2$, $\tilde{L}_2$, $\tilde{E}_2$, and $\tilde{C}_2$ by repeated application of Lemma 3.1.6. For C_2 , choose any chamber attached to R_2 other than $\tilde{C}_2$. This ensures that γ_1 is no bigger than intended, and that $\gamma_2 = 0$. Choose the rest of P arbitrarily.

Subcase 3: $\gamma_1 + T(P_b) - 1 < \gamma_1 + \gamma_2$, and point a in Figure 3.64 is at least as far right as point d (the alternative is Figure 3.65, which we will consider in subcase 4). Point a is a distance of $\gamma_1 + \gamma_2 - T(P_b) + 1$ edge lengths from the right boundary of P , and point d is a distance of $\gamma_1 + T(P_b) - 1$ edge lengths from the same boundary. So our condition on a and d is the same as saying $\gamma_1 + \gamma_2 - T(P_b) + 1 \leq \gamma_1 + T(P_b) - 1$, i.e., $\gamma_2 + 2 \leq 2T(P_b)$. Choose Q_1 and $\tilde{r}_1$ from $b\sigma(A_1^{\frac{1}{2}})$. Let $R_1 = \sigma^{-1}b^{-1}Q_1$. Then $R_1 \subseteq A_1^{\frac{1}{2}}$, so L_1 is in A_M , and Q_1 and $\tilde{r}_1$ are attached to $\overline{P}$ along $b\sigma(K)$. Now consider the apartment made up of $A_1^{\frac{1}{2}}$, $b\sigma(A_1^{\frac{1}{2}})$, and the strip in A_M between K and $b\sigma(K)$. Extend the line l in Figure 3.64 to a wall in this apartment, and take $A_2^{\frac{1}{2}}$ to be a half apartment coming out of this wall. Choose Q_2 , $\tilde{r}_2$, L_2 and N_1 from $A_2^{\frac{1}{2}}$. This determines R_2 . Choose Q_3 arbitrarily. This determines $\tilde{r}_3$, $\tilde{L}_3$ and $\tilde{N}_3$, none of which will be part of the final construction. Choose N_2 using Lemma 3.1.6. Let $\tilde{C}$ and $\tilde{D}$ be the unique chambers attached to $\tilde{N}_3$ and N_2 , given by Lemma 3.1.6. Choose $C \neq \tilde{C}$. This assures that γ_2 is no bigger than desired. Construct the rest of P arbitrarily.

Subcase 4: $\gamma_1 + T(P_b) - 1 < \gamma_1 + \gamma_2$, and point a in Figure 3.65 is strictly to the left of point d (so $\gamma_2 + 2 > 2T(P_b)$ and the condition $\gamma_1 + T(P_b) - 1 < \gamma_1 + \gamma_2$ become redundant). Construct R_1 , L_1 , Q_1 and $\tilde{r}_1$ exactly as in subcases 3 and 1. Construct N_1 , L_2 , $\tilde{r}_2$, and Q_2 exactly as in subcase 3. This determines R_2 and r_1 . Choose Q_3




Figure 3.62: Existence of all values of γ_1 and γ_2 , case 2 subcase 1




Figure 3.63: Existence of all values of γ_1 and γ_2 , case 2 subcase 2




Figure 3.64: Existence of all values of γ_1 and γ_2 , case 2 subcase 3

arbitrarily. This determines R_3 . Use Lemma 3.1.6 repeatedly to fill in $\tilde{r}_3$, L_3 and N_2 . Choose the unique possible C_3 and D_3 to connect N_2 and L_3 . Repeat this process to point a , then, at the next step choose for C_i any chamber other than the unique one that connects to L_i . This ensures γ_2 is no bigger than desired. Choose the rest of P arbitrarily.

Up to this point, we have defined $\overline{P}$, P_σ , and the invariants γ_1 and γ_2 for $b = 1$ or b with $\alpha + 2\beta \neq 0$, and $(t, e) \in I_1$. We have also shown that if $1 \leq \gamma_1 \leq \gamma_1^{\max}$, $0 \leq \gamma_2 \leq \gamma_2^{\max}(\gamma_1)$, then there is some $x \in SL_3(L)$ such that xC_M gives rise to (t, e) , γ_1 and γ_2 . We now broaden the definition of γ_2 to include negative values. We will then have to re-address the issue of whether all possible γ_2 actually occur. Consider Figure 3.66. If there is no chamber connecting chamber 7 to chamber 5, we say $\gamma_2 \leq 0$ (previously we just said $\gamma_2 = 0$). In this case we ask if there is a chamber connecting 4 to 8. If not, then $\gamma_2 = 0$. If so, then $\gamma_2 \leq -1$, and we call the connecting chamber $\tilde{7}$. In this case, we fill in chambers $\tilde{6}$ and $\tilde{9}$ using chambers $\tilde{7}$, 8, 11, 10 and Lemma 3.1.6. We now ask if there is a chamber connecting $\tilde{6}$ to 2. If not, then $\gamma_2 = -1$, and if so then $\gamma_2 \leq -2$. We proceed in this way to determine the value of γ_2 . Note that if $\gamma_2^{\min}(\gamma_1) = -\min\{S(\overline{P}), T(P) - \gamma_1\}$, then $\gamma_2 \geq \gamma_2^{\min}$ automatically.

We now show that given any b such that $b = 1$ or $\alpha + 2\beta \neq 0$, given $(t, e) \in I_1$, and given γ_1 and γ_2 such that $1 \leq \gamma_1 \leq \gamma_1^{\max}$ and $\gamma_2^{\min}(\gamma_1) \leq \gamma_2 \leq \gamma_2^{\max}(\gamma_1)$, there exists an x such that x and b give rise to (t, e) , γ_1 and γ_2 . Given what we have already done, we may assume that $\gamma_2 \leq 0$. As usual we proceed in cases.

Case 1: $b = 1$. See Figure 3.67. Let L be a wall in A_M , and let $A_1^{\frac{1}{2}}$ be a half apartment in $\mathcal{B}_2$ coming out of L . Choose Q_1 in $A_1^{\frac{1}{2}}$. Choose chamber 4 in $\mathcal{B}_6 \setminus \mathcal{B}_2$. This determines 8, and therefore by Lemma 3.1.6 also determines $\tilde{3}$ and $\tilde{7}$ connecting 4 and 8. To arrange $\gamma_2 = 0$, choose 7 such that $7 \neq \tilde{7}$ and $\sigma(7) \neq \tilde{3}$, and construct the rest of P arbitrarily. To arrange $\gamma_2 \leq -1$, choose $7 = \sigma^{-1}(\tilde{3})$. Since $4, 8 \in \mathcal{B}_6 \setminus \mathcal{B}_2$, $\sigma(\tilde{7}) \neq \tilde{3}$, so $\tilde{7} \neq 7$, so $\gamma_2 \leq 0$ still holds. On the other hand $3 = \sigma(7) = \tilde{3}$, so 3 can be connected to 8 (via $\tilde{7}$), ensuring $\gamma_2 \leq -1$. Choose $\tilde{6}$ and $\tilde{9}$ to fit between $\tilde{7}$ and 10 using Lemma 3.1.6. Choose $\tilde{1}$ and $\tilde{5}$ to fit between 2 and $\tilde{6}$. To arrange $\gamma_2 = -1$ choose $1 \neq \tilde{1}$ and construct the rest of P arbitrarily. To arrange $\gamma_2 \leq -2$, choose $1 = \tilde{1}$. Continue in this manner.

Case 2: $b \neq 1$. We break this case into subcases.

Subcase 1: $T(P_b) = 1$. See Figure 3.68. We let K and K_1 be walls in A_M , K a horizontal wall, and K_1 a wall such that $bK_1 = K_1$. Let $\tilde{T}$ be the first row of chambers in a half apartment $A_1^{\frac{1}{2}}$ coming out of K_1 to the left. Choose $A_1^{\frac{1}{2}}$ such that $b\sigma(\tilde{T}) \neq \tilde{T}$. This implies that $b\sigma(\tilde{T})$ has no chambers in common with $\tilde{T}$. Let A be the apartment made up of A_M on the right of K_1 and $A_1^{\frac{1}{2}}$ on the left. Then half of K is in A , and we can extend this half to $\tilde{K}$, a wall in A . Let $A_2^{\frac{1}{2}}$ be a half





Figure 3.65: Existence of all values of γ_1 and γ_2 , case 2 subcase 4

Figure 3.66: Definition of negative values of γ_2




Figure 3.67: Existence of all negative values of γ_2 , case 1

apartment coming out of $\tilde{K}$, and choose R_1, R_2 and $\tilde{C}_2$ from $A_2^{\frac{1}{2}}$. We can now throw away all of $A_1^{\frac{1}{2}}$ except $\tilde{T}$. We get $Q_1 = b\sigma(R_1)$, $Q_2 = b\sigma(R_2)$, P_b , and $L_1 \subseteq A_M$. Note that $b\sigma(\tilde{T})$ is another row of chambers coming out of L , and note that we have $\tilde{D}_2 = b\sigma(\tilde{C}_2)$ attaching Q_2 to $b\sigma(\tilde{T})$, as drawn in the figure. We have $\tilde{F}_2 \subseteq b\sigma(\tilde{T})$ which connects $\tilde{D}_2$ to L_1 , and by Lemma 3.1.6, we have $\tilde{E}_2$ and $\tilde{C}_2$ attaching $b\sigma(\tilde{T})$ to R_2 as drawn. Note that $\tilde{\tilde{C}}_2 \neq \tilde{C}_2$. If these chambers were equal, we could create a contradiction to the uniqueness part of Lemma 3.1.6 by producing two galleries of length two from the edge between $\tilde{\tilde{C}}_2 = \tilde{C}_2$ and $\tilde{E}_2$, down through $\tilde{T}$ and $b\sigma(\tilde{T})$ respectively, to L_1 . To make sure $\gamma_2 \leq 0$, choose $C_2 \neq \tilde{\tilde{C}}_2$. To make sure $\gamma_2 \leq -1$, choose $C_2 = \tilde{\tilde{C}}_2$. For $\gamma_2 = 0$, choose any other C_2 . If we chose $\gamma_2 = 0$, fill out the rest of P arbitrarily. If we chose $\gamma_2 \leq -1$, then what we have done becomes the base case of an induction.

Let $D_2 = b\sigma(C_2)$, and see Figure 3.69. The region made up of $\tilde{L}_2, \tilde{S}_2$ is taken from $b\sigma(\tilde{T})$. We assume by induction that the $Q_j, D_j, \tilde{L}_j$ and $\tilde{S}_j$ for $2 \leq j \leq i$ are all contained in a single half apartment, and that $\gamma_2 \leq -i + 1$. We choose Q_{i+1} arbitrarily. If we want $\gamma_2 = -i + 1$, we choose D_{i+1} such that there is no chamber attaching it to $\tilde{L}_i$. If we want $\gamma_2 \leq -i$, we choose a chamber D_{i+1} that can be attached to $\tilde{L}_i$. This enables us to fill in an $\tilde{L}_{i+1}$ and an $\tilde{S}_{i+1}$ uniquely using Lemma 3.1.6. We can continue thus until we either reach the desired value for γ_2 , or until we are stopped by $\gamma_2^{\min}(\gamma_1)$.

Subcase 2: $T(P_b) > 1$. See Figure 3.70. Construct R_1 and Q_1 in the standard way. Choose Q_2 such that there is no chamber connecting it to L_1 (to ensure $\gamma_2 \leq 0$). If we want $\gamma_2 = 0$, we choose D_2 such that there is no chamber connecting it to the line l (which is in L_1 and P). If we want $\gamma_2 \leq -1$, we choose D_2 such that there is a chamber $\tilde{E}_2$ connecting it to the line l . Note that $\tilde{E}_2 \not\subseteq L_1$. Construct $\tilde{S}_2$ using Lemma 3.1.6. The region $\tilde{S}_2$ does not have any chambers in common with L_1 or with P . We then choose Q_3 arbitrarily, and if we want $\gamma_2 = -1$, we choose D_3 such that there is no chamber connecting it to S_2 . If we want $\gamma_2 \leq -2$, we choose D_3 such that there is such a chamber. Proceed in this way.

We have now shown that given a σ -conjugacy class b such that either $b = 1$ or $\alpha + 2\beta \neq 0$, and given a type edge pair $(t, e) \in I_1$, one can find $x \in SL_3(L)$ such that xC_M has SMG of type t and departure edge e , and such that the invariants γ_1, γ_2 associated with the composite gallery Γ_x are as desired (within the limitations $1 \leq \gamma_1 \leq \gamma_1^{\max}$ and $\gamma_2^{\min}(\gamma_1) \leq \gamma_2 \leq \gamma_2^{\max}(\gamma_1)$). This is useful because we will now prove that $\rho(x^{-1}b\sigma(x)C_M)$ is determined by b, t, e, γ_1 , and γ_2 . Again, we consider cases.

Case 1: $\gamma_2 > 0$. See Figure 3.71, which represents the structure $\overline{P} \cup P_\sigma$ transported back to C_M (in other words, $x^{-1}(\overline{P} \cup P_\sigma)$). We can find $g \in I$ that sends the shaded part of this picture to A_M . This is because one can find an apartment containing the shaded part of the figure but not containing the unshaded part. Therefore g does




Figure 3.68: Existence of all negative values of γ_2 , case 2 subcase 1




Figure 3.69: Existence of all negative values of γ_2 , case 2 subcase 1 diagram 2




Figure 3.70: Existence of all negative values of γ_2 , case 2 subcase 2




Figure 3.71: $x^{-1}(\overline{P} \cup P_\sigma)$ for $\gamma_2 > 0$

not send the unshaded part of the diagram to A_M . So we assume from this point forward that the shaded part of Figure 3.71 is in A_M and the unshaded part is not. It is easy to see now that ρ has the effect of reflecting the chambers in R_1 across l_1 , reflecting the chambers in R_2 across l_2 , reflecting the chambers in R_3 first across l_1 and then across l_3 , and reflecting the chambers in R_4 across l_2 . So one can determine the value of $\rho(x^{-1}b\sigma(x)C_M)$. A specific example is given in Figure 3.72. Note that the existence of a corner at point x in the figure is necessary for the considerations that arrive at these results.

Case 2: $\gamma_2 \leq 0$. See Figure 3.73. Once again, we can arrange for the shaded part of the figure to be in A_M , and for the unshaded part to not be in A_M . According to the value of γ_2 , we can connect P_σ to l using region $\tilde{R}_1$ in Figure 3.74. Here, l is in $\overline{P}$, and $\tilde{R}_1$ shares no chambers in common with $\overline{P}$. We can now fill in $\tilde{R}_2$ uniquely using Lemma 3.1.6 so that it is connected above to P_σ and on the right to $\tilde{R}_1$. There now exist chambers $\tilde{C}$ and $\tilde{D}$ such that neither is in $\overline{P}$, but such that $\tilde{C}$ is connected to




Figure 3.72: Example of how γ_1 and γ_2 determine $\rho(x^{-1}b\sigma(x)C_M)$

$\tilde{R}_2$, and $\tilde{D}$ has its right edge in $\overline{P}$. Also, $\tilde{C}$ and $\tilde{D}$ are connected to each other. We fill in $\tilde{R}_3$ using Lemma 3.1.6. With respect to folding, the corner labeled with an x is similar in function to the corner labeled with an x in Figure 3.71. As such, referring to Figure 3.75, chamber $\tilde{D}$ is reflected across l_2 by ρ , the chambers in region $\tilde{B}_1$ (some of which are in P_σ and some of which are not in $\overline{P} \cup P_\sigma$) are reflected across l_1 , the chambers in $\tilde{B}_2$ (some of which are in P_σ and some of which are not in $\overline{P} \cup P_\sigma$) are reflected first across l_1 , then across l_3 , and the chambers in $\tilde{B}_3$ are reflected across l_2 . So one can determine the value of $\rho(x^{-1}b\sigma(x)C_M)$ for any specific example.

In Section 3.1.2, we computed chambers to include in the superset S_1 by computing all possible foldings of the composite gallery Γ_x for x giving rise to any fixed type-edge pair (t, e) . The set S_1 consisted of the union of these results for all (t, e) in $I_1 \cup I_2$. We are now in a position to show that for all $(t, e) \in I_1$, the collection of possible foldings of Γ_x gives rise only to chambers in S . This, combined with similar results for I_2 proves that $S = S_1$ for $b = 1$ or b with $\alpha + 2\beta \neq 0$.

Let $S_{(t,e)}^f$ be the collection of chambers obtained by computing the possible foldings of Γ_x for x giving rise to the type-edge pair (t, e) . So

$$S_1 = \left[\bigcup_{(t,e) \in I_1} S_{(t,e)}^f \right] \cup \left[\bigcup_{(t,e) \in I_2} S_{(t,e)}^f \right].$$




Figure 3.73: $x^{-1}(\overline{P} \cup P_\sigma)$ for $\gamma_2 \leq 0$




Figure 3.74: $x^{-1}(\overline{P} \cup P_\sigma)$ for $\gamma_2 \leq 0$, folding explained




Figure 3.75: $x^{-1}(\overline{P} \cup P_\sigma)$ for $\gamma_2 \leq 0$, folding explained further




Figure 3.77: A composite gallery for $p = 1$, $\alpha = 2$, $\beta = -1$

of $\rho(x^{-1}b\sigma(x)C_M)$ is in $S_{(t,e)}^\gamma$ because we may choose γ_1 and γ_2 such that $\gamma_1 + \max\{\gamma_2, 0\} = S(P_\sigma) + i - 1$ (to ensure that the first change choice occurs at d_i) and such that $\gamma_1 - \min\{0, \gamma_2\} = \frac{s+1}{2}$ (to ensure that the second change choice occurs s edges after d_i). If no second change choice is desired, then choose $\gamma_2 = \gamma_2^{\min}(\gamma_1)$.

If one makes the first change choice at some e_i , then only one subsequent change choice is possible. If such a change choice is made, it must be done at one of the u_j . The resulting $\rho(x^{-1}b\sigma(x)C_M)$ is in $S_{(t,e)}^\gamma$ because we may choose γ_1 and γ_2 such that $\gamma_1 + \max\{\gamma_2, 0\} = i$ and such that $\gamma_1 - \min\{\gamma_2, 0\} = j$. If no second change choice is desired, choose $\gamma_2 = \gamma_2^{\min}(\gamma_1)$.

In essence, the choice of γ_1 and γ_2 above corresponds to choosing the point x in Figures 3.71 and 3.75 such that l_1 passes through the first change choice and l_2 passes through the second.

We now address the situation where $(t, e) \in I_1$ still, but now $b \neq 1$ and $\alpha + 2\beta = 0$. As before, let p be the number of chambers in Γ_E^3 between the edge of departure and the turning edge. So $p \geq 1$ is odd. We first consider the case where $p = 1$. So, fixing t, e, b with the specified properties, and choosing $y \in SL_3(L)$ such that y gives rise to (t, e) , and given some possible folding of a gallery of the same type as $y^{-1}\Gamma_y$, we want to show that there is $x \in SL_3(L)$ such that $\rho(x^{-1}b\sigma(x)C_M)$ is the last chamber of this folding.

For $p = 1$, a sample composite gallery for b with $\alpha = 2$, $\beta = -1$ is pictured in Figure 3.77. The choice points are marked. For general p, b , we get a similar structure, with every edge in Γ_E^3 a choice point, and the edge between Γ_E^2 and Γ_E^3 another choice point. Note that for $p = 1$, if one makes a change choice at any choice point other than the horizontal one between Γ_E^2 and Γ_E^3 , no subsequent choices are available. A change choice at the horizontal change point allows at most two more choices.

Let K be a wall in A_M such that $bK = K$. Let T be the first row of chambers in some half apartment coming out of K . Choose $T \subseteq \mathcal{B}_1$, so σ does not affect T . Then $b\sigma(T) = T$. Choose the first i chambers of Γ_E^3 in T , and the rest not in




Figure 3.78: All foldings of composite galleries with $p = 1$ occur, where $b \neq 1$ and $\alpha + 2\beta = 0$

T . So $\Gamma_E^1 = \sigma^{-1}b^{-1}(\Gamma_E^3)$ and $\Gamma_E^2 \subseteq A_M$ are determined. This composite gallery construction folds in such a way that the first (and only) change choice occurs in Γ_E^3 between the i th and $(i+1)$ st chambers. To arrange no change choices at all, choose $\Gamma_E^3 \subseteq T$.


To arrange that the edge between Γ_E^2 and Γ_E^3 be a change choice, take $T \subseteq \mathcal{B}_2 \setminus \mathcal{B}_1$. Then $b\sigma(T) \cap T = K$. If we want no subsequent change choice, then choose $\Gamma_E^3 \subseteq b\sigma(T)$. If we want the second change choice to be between the i th and $(i+1)$ st chambers in Γ_E^3 , then choose the first i chambers of Γ_E^3 in $b\sigma(T)$, and the rest outside of $b\sigma(T)$. We now have a situation for which Figure 3.78 is an example (here b has $\alpha = 4$ and $\beta = -2$). The galleries Γ_E^1 and Γ_E^3 are drawn as part of T and $b\sigma(T)$, respectively, in this diagram, even though only part of each need be.

Let e be the edge in Γ_E^3 at which the second change choice is to occur. Then the chambers of Γ_E^3 that are part of $b\sigma(T)$ are exactly those that occur before e . If e is an edge of the same angle as e_1 in Figure 3.78, then no subsequent change choice is possible. If e is of the same angle as e_2 , then e_1 could be a third (and final) change choice. To arrange this, choose Γ_E^3 in such a way that there is no chamber connecting edge g to edge h . To arrange no third change choice, choose Γ_E^3 such that there is a chamber connecting g to h .

We have now shown for $p = 1$ that any possible folding can occur. We now assume inductively that any possible folding can occur for $p \leq n$. Suppose that x gives rise to a type-edge pair (t, e) with $p = n + 1$, and a composite gallery Γ_x . We begin by defining an invariant γ_3 of x . Consider Figure 3.79, which shows an example composite gallery for b with $\alpha = 2$, $\beta = -1$. We use Lemma 3.1.6 to add vertices v_1 and v_2 to the structure. Note that v_2 may or may not be part of Γ_x . If v_2 is in Γ_x then we say $\gamma_3 \geq 1$. If v_2 is not in Γ_x , then we say $\gamma_3 = 0$. Although we have only given the definition of γ_3 for the specific example of Figure 3.79, it is clear how one would proceed in general. Note that γ_3 does not necessarily have a specific value. We either have $\gamma_3 = 0$ or $\gamma_3 \geq 1$.

In case $\gamma_3 \geq 1$, we can find a gallery from xC_M to $b\sigma(x)C_M$ that is of the same shape as a composite gallery of some type-edge pair with $p = n$. Therefore, by induction, we need not worry about this case.

If $\gamma_3 = 0$, then the edge between Γ_E^2 and Γ_E^3 is forced to be a change choice. For instance, the gallery in Figure 3.79 becomes the gallery in Figure 3.80 upon addition of the vertex v_3 (which can be done using Lemma 3.1.6). We can use techniques

Figure 3.79: Definition of γ_3

similar to those used in the first part of this section to show that one can arrange for such a gallery to fold in any desired way.

To finish the proof that $S_1 = S$, it would be necessary to consider $(t, e) \in I_2$. One should consider the case $b = 1$ and the cases $2\alpha + \beta \neq \alpha + 2\beta$ separately from the cases in which $b \neq 1$ but $2\alpha + \beta = \alpha + 2\beta$. The process is similar to what we have already done, and is omitted.

3.1.7 Symmetry Under a $\mathbb{Z}/3$ -action

In Section 2.1.3 we noticed from results developed in Sections 2.1.1 and 2.1.2 that for SL_2 , the set of w with non-empty $X_w(b\sigma)$ has $\mathbb{Z}/2$ -symmetry about the center of C_M . We gave an *a priori* proof of this fact. The results developed for SL_3 in the previous sections of this chapter indicate that in this case the set of w with non-empty $X_w(b\sigma)$ has $\mathbb{Z}/3$ -rotational symmetry about the center of C_M . We adapt the proof from Section 2.1.3 to give an *a priori* proof of the SL_3 symmetry.

Let

$$q = \begin{pmatrix} 0 & 0 & \pi \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \in GL_3(F).$$

This matrix acts on A_M by rotating it 120° counterclockwise about the center of C_M . Let

$$g = \begin{pmatrix} \pi & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \in GL_3(F).$$

considered as elements of GL_3 or PGL_3 . Further, any two σ -conjugacy classes of SL_3 remain distinct when we pass to σ -conjugacy classes of GL_3 or PGL_3 . It is also true that GL_3 and PGL_3 have more σ -conjugacy classes than SL_3 , just as was the case for the analogous rank one groups. Unlike in the rank one case, we will not address these additional σ -conjugacy classes, since for one thing we have not even addressed all the σ -conjugacy classes of SL_3 itself.

If b is one of the matrices listed at the beginning of Section 3.1, this time considered as an element of GL_3 or PGL_3 , then we ask for a description of $\{inv(x, b\sigma(x)) : x \in G(L)\}$, i.e., we want to describe $\{(\rho(x^{-1}b\sigma(x)C_M), v(\det(x^{-1}b\sigma(x)))) \in W_a \ltimes M : x \in G(L)\}$. We note that $v(\det(x^{-1}b\sigma(x))) = v(\det(b))$ is fixed for fixed b (and equal to 0 for the b we have chosen). So we need to examine $\{\rho(x^{-1}b\sigma(x)C_M) \in W_a : x \in G(L)\}$, which, *a priori*, may be bigger than $\{\rho(x^{-1}b\sigma(x)C_M) \in W_a : x \in SL_3(L)\}$, since $G(L)$ acts on $\mathcal{B}_\infty$ in ways that SL_3 does not. However, if q is as defined in Section 3.1.7, and if $y = q^{-v(\det(x))}$ then $v(\det(xy)) = 0$. Note that if $G = PGL_3$, then $v(\det(x))$ is only determined mod 3. But since q^3 is a scalar matrix, this does not cause problems with the definition of y . Now $\det(xy) \in \mathcal{O}_L^\times$, and since $(\mathcal{O}_L^\times)^3 = \mathcal{O}_L^\times$, we may further modify xy on the right by a scalar matrix of determinant $\det(xy)^{-1}$. Therefore we may assume without loss of generality that $\det(xy) = 1$, i.e., $xy \in SL_3(L)$. But $xyC_M = xC_M$, so $\sigma(xy)C_M = \sigma(x)C_M$, so $b\sigma(xy)C_M = b\sigma(x)C_M$, so $x^{-1}b\sigma(xy)C_M = x^{-1}b\sigma(x)C_M$, so $\rho(x^{-1}b\sigma(xy)C_M) = \rho(x^{-1}b\sigma(x)C_M)$. But it is easy to see that $\rho(y^{-1}x^{-1}b\sigma(xy)C_M)$ is just a rotation of $\rho(x^{-1}b\sigma(xy)C_M)$ about the center of C_M by $(120^\circ)(v(\det(x)))$ counterclockwise. Since we already proved that the set $\{\rho(x^{-1}b\sigma(x)C_M) : x \in SL_3(L)\}$ is rotation invariant by 120° and 240° , we see that $\{\rho(x^{-1}b\sigma(x)C_M) : x \in SL_3(L)\} = \{\rho(x^{-1}b\sigma(x)C_M) : x \in G(L)\}$.

3.3 Sp_4

We have applied some of the methods of Section 3.1 to Sp_4 . This section contains outlines of the application of these methods, and the results. The structure of the section is the same as that of Section 3.1.

The version of $Sp_4(E)$ that we will use for E/F a field extension is the fixed points in $GL_4(E)$ of the involution $\Theta : GL_4(E) \rightarrow GL_4(E)$ where $\Theta(g) = M^{-1}(g^t)^{-1}M$ for

$$M = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}.$$

Note that if T is the standard maximal torus in GL_4 , then the fixed points T^Θ form a split maximal torus in Sp_4 , and if B is the standard Borel subgroup of GL_4 , then the fixed points B^Θ form a Borel for Sp_4 .




Figure 3.81: How the torus of Sp_4 acts on A_M

The letter b in this section will always stand for an element of Sp_4 of the form

$$\begin{pmatrix} \pi^\alpha & 0 & 0 & 0 \\ 0 & \pi^\beta & 0 & 0 \\ 0 & 0 & \pi^{-\beta} & 0 \\ 0 & 0 & 0 & \pi^{-\alpha} \end{pmatrix},$$

where $\alpha \geq \beta \geq 0$. These represent distinct σ -conjugacy classes in $Sp_4(L)$, but they do not constitute a complete collection of σ -conjugacy classes. [6] [7]

To understand something about how Sp_4 acts on the main apartment of its building, consider Figure 3.81. The main chamber and the main vertex are labeled in this figure, and the value of bC_M with $\alpha = 1$ and $\beta = 0$ is labeled with an α . The value of bC_M with $\alpha = 0$ and $\beta = 1$ is labeled with a β .

3.3.1 Standard Minimal Galleries and Composite Galleries

We define the four *primary directions* D_1, D_2, D_3, D_4 and the four *secondary directions* d_1, d_2, d_3, d_4 in A_M for Sp_4 as marked in Figure 3.82. Given a chamber $E \subseteq A_M$, we define the *standard minimal gallery* (SMG) between C_M and E as follows. If E is in one of the corridors marked $c_1, \dots, c_8$ on Figure 3.82, then the SMG is the unique minimal gallery from C_M to E . If E is in region R_i , proceed first in direction D_i , then in direction d_i . If E is in region r_i , proceed first in direction D_i , then in direction d_j , where $j \equiv i + 1 \pmod{4}$.

As was the case for SL_3 , one can prove that if E is now allowed to be a chamber anywhere in the building $\mathcal{B}_\infty$ of $Sp_4(L)$, then there exists a unique gallery G_E between




Figure 3.82: Primary and Secondary Directions in Sp_4

C_M and E which is minimal and such that $\rho(G_E)$ is the SMG from C_M to $\rho(E)$. We define the SMG from C_M to E to be G_E in this case.

We construct the *composite gallery* Γ_E associated to E as in the SL_3 case. We also define the *edge of departure* in the same way as for SL_3 .

3.3.2 A Conjectural Superset of the Solution Set

Just as for SL_3 , we can arrive at a superset S_1 of the solution set S by computing $S_1 = \cup_{(t,e)} S_{(t,e)}$, where the union is over all type-edge pairs (t, e) , and $S_{(t,e)}$ is the collection of final chambers of possible foldings of composite galleries arising from SMGs of type t and edge of departure e . This computation is even more prohibitively lengthy than was the case for SL_3 , and has only been carried out to completion for $b = 1$, for which the results are in Figure 3.83.

Let I_1 be the infinite class of type-edge pairs (t, e) with t the type of the SMG of some chamber in r_2 , and e some horizontal departure edge. Let I_2 be the infinite class of type-edge pairs (t, e) with t the type of the SMG of some chamber in c_4 , and e an arbitrary departure edge. These I_1 and I_2 are analogous to the infinite classes of the same name for SL_3 .

Conjecture 3.3.1. $S_1 = \cup_{(t,e)} S_{(t,e)}$, where the union is over all type-edge pairs in $I_1 \cup I_2$.

Computation of $\cup_{(t,e)} S_{(t,e)}$ as (t, e) ranges over $I_1 \cup I_2$ is a lengthy but reasonable computation. The details of this computation for any b are discussed in Section 3.3.3. For $\alpha = 3$ and $\beta = 1$, the results are given in Figure 3.84. For $\alpha = 6$, $\beta = 3$, the results are given in Figure 3.85, for $\alpha = 6$, $\beta = 5$ the results are given in Figure 3.86, and for $\alpha = 7$, $\beta = 1$ the results are given in Figure 3.87. The lines on these figures and on Figure 3.83 are only there to make the figures easier to look at. The chambers in Figure 3.84 that are shaded more darkly are the chambers $w^{-1}bwC_M$ for $w \in W$.

The first piece of evidence supporting Conjecture 3.3.1 is that for $\alpha = \beta = 0$, I_1 and I_2 yield Figure 3.83, which is the complete computation $\cup_{(t,e)} S_{(t,e)}$, as (t, e) ranges over all type-edge pairs.

The chambers marked with a $*$ in Figures 3.84, 3.85, 3.86, and 3.87 provide evidence against Conjecture 3.3.1, as they seem to be holes in the pattern. However, it could be the case that these chambers are actually missing from the true $S_1 = \cup_{(t,e)} S_{(t,e)}$, where the union is over all (t, e) . Preliminary computations not presented seem to provide some evidence that this is the case, in support of Conjecture 3.3.1. However, this evidence is not a strong indication.




Figure 3.83: Result for $b = 1$, Sp_4




Figure 3.84: Result for $\alpha = 3$, $\beta = 1$, Sp_4




Figure 3.85: Result for $\alpha = 6$, $\beta = 3$, Sp_4




Figure 3.86: Result for $\alpha = 6$, $\beta = 5$, Sp_4




Figure 3.87: Result for $\alpha = 7$, $\beta = 1$, Sp_4

3.3.3 Relationship Between $X_w(1\sigma)$ and $X_w(b\sigma)$ for Sp_4 , and an Efficient Way of Computing Supersets

In Section 3.1.3 we gave a more efficient method of computing the results of the folding of $I_1 \cup I_2$. In that case, we knew these results gave the entire superset S_1 of the solution set. One can apply the methodology of Section 3.1.3 to Sp_4 to produce an efficient way of computing the folding of $I_1 \cup I_2$ for Sp_4 . However, we have only conjectured that the folding of $I_1 \cup I_2$ for Sp_4 gives the entire superset S_1 .

Since the methods of this section are very similar to those of Section 3.1.3, we present only results.

Proposition 3.3.1. For b with $\alpha > \beta$, computing the folding results of $I_1 \cup I_2$ is the same as computing the folding results of some collection of half-infinite galleries analogous to those pictured in Figure 3.88, and combining those results with their own reflection across the vertical line of symmetry of C_M . For b with $\alpha = \beta$, computing the folding results of $I_1 \cup I_2$ is the same as computing the folding results of some collection of half-infinite galleries analogous to those pictured in Figure 3.89, and combining these results with their reflection across the vertical line of symmetry of C_M . Although only one specific value of b is pictured in each figure, it is clear what the situation would be for arbitrary b . In other words, it is clear what the analogous collection of half-infinite galleries would be for any b .

3.3.4 The Method of Kottwitz and Rapoport Applied to Sp_4

As in the SL_3 case, we produce a subset $S_2 \subseteq S$ for $b = 1$. Let $a \in Sp_4(F)$ be of the form

$$a = \begin{pmatrix} \pi^m & 0 & 0 & 0 \\ 0 & \pi^n & 0 & 0 \\ 0 & 0 & \pi^{-n} & 0 \\ 0 & 0 & 0 & \pi^{-m} \end{pmatrix}$$

where there are no conditions on m and n . Let w be one of the following matrices:

$$r = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 \end{pmatrix} \quad ; \quad r^2 = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \quad ; \quad r^3 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Note that $r^4 = -1$. One can check that in fact $r \in Sp_4(F)$ by computing $\Theta(r) = r$. The r^i are representatives of Weyl group elements for Sp_4 . In fact, $W = D_4$, and r is a representative of a generator of the order 4 cyclic subgroup of W .

The matrix aw belongs to a basic σ -conjugacy class if there is some l such that $aw\sigma(aw)\sigma^2(aw)\cdots\sigma^{l-1}(aw)$ is central in Sp_4 . Since Sp_4 is simply connected, $b = 1$ is the only basic σ -conjugacy class. [6] [7]




Figure 3.88: Half infinite galleries for Sp_4 , $\alpha > \beta$ (specifically, $\alpha = 3$, $\beta = 1$)




Figure 3.89: Half infinite galleries for Sp_4 , $\alpha = \beta$ (specifically, $\alpha = \beta = 3$)




Figure 3.90: Main results of the method suggested by Rapoport and Kottwitz, Sp_4

Lemma 3.3.2. There exists an l such that $aw\sigma(aw)\sigma^2(aw)\cdots\sigma^{l-1}(aw)$ is central for any choice of m and n .

Proof. This is similar to Lemma 3.1.3. The only difference is that $l = 4$. $\square$

Therefore, the double- I -cosets pictured in Figure 3.90 all meet the σ -conjugacy class of 1 non-trivially.

If w is a representative of an element of W other than r^i for $i = 1, 2, 3$, and if certain conditions are placed on m and n , then it is still possible for $(aw)^l$ to be central for some l .

Lemma 3.3.3. If $a = 1$, then for $l = 4$, $(aw)^l$ is central for any $w \in W$.




Figure 3.91: The $a = 1$ results of the method suggested by Rapoport and Kottwitz, Sp_4

This means the double- I -cosets corresponding to chambers shaded in Figure 3.91 all intersect the σ -conjugacy class of $b = 1$ non-trivially. Let

$$f = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Lemma 3.3.4. If $n = 0$ then $aa^w a^{w^2} a^{w^3} w^4 = 1$ for $w = rf$. If $m = 0$, then $aa^w a^{w^2} a^{w^3} w^4 = 1$ for $w = r^3 f$. If $m = -n$, then $aa^w a^{w^2} a^{w^3} w^4 = 1$ for $w = f$. If $m = n$, then $aa^w a^{w^2} a^{w^3} w^4 = 1$ for $w = r^2 f$.

Proof. Compute the relevant matrix products. □

The implication of this lemma is that the double- I -cosets corresponding to the chambers in Figure 3.92 all intersect the σ -conjugacy class of 1 non-trivially.

Just as the methods of Section 3.1.4 generalize to SL_n , the results of this section generalize to Sp_{2n} .

3.3.5 A Subset of the Solution Set for $b = 1$ for Sp_4

This section is the Sp_4 analogue of Section 3.1.5, and the setup is the same. In slightly different language, we choose a gallery G of length 2 or 3 starting at C_M and having only its first chamber C_M in A_M . We denote the chambers of G after C_M by $G_1, \dots, G_i$ ($i = 1$ or 2), and we let e be the edge by which G_1 is adjacent to C_M . We let Γ^2 be a minimal gallery from e to $\tilde{b}e$, where $\tilde{b}C_M$ is one of the chambers in A_M obtained in the previous section. We form a composite gallery Γ from $G_1, \dots, G_i, \Gamma^2$, and $\tilde{b}\sigma(G_1), \dots, \tilde{b}\sigma(G_i)$. The collection of possible foldings of $x^{-1}\Gamma$ gives candidates




Figure 3.92: Results of the method suggested by Rapoport and Kottwitz under the circumstances of Lemma 3.3.4, Sp_4




Figure 3.93: The choices D_1 , D_2 and D_3 of G_1

for an addition to S_2 . If there is only one possible folding then it gives us a new element of S_2 .

Starting with $i = 1$ and $xC_M = G_1$ equal to one of the three chambers D_1 , D_2 , D_3 pictured in Figure 3.93, we get results as pictured in Figures 3.94, 3.95, and 3.96 for $\tilde{b} = aw$ with $w = r$, Figures 3.97, 3.98, and 3.99 for $w = r^2$, and Figures 3.94, 3.95, and 3.96 again for $w = r^3$. Occasionally during these computations, the composite gallery Γ is not minimal, and so $\rho(x^{-1}\Gamma)$ cannot be determined (although in all such cases $\rho(x^{-1}\Gamma)$ has one of two possible values). In these cases, neither of the possible values of $\rho(x^{-1}\Gamma)$ is included in the result figures. So all chambers in figures 3.94 through 3.99 are actually in S_2 .

Combining the results obtained so far in this section with those of the previous section gives Figure 3.83, which was already known to be the superset $S_1 \supseteq S$. Therefore there is no need to consider the $i = 2$ case, and we see that $S_2 = S_1$, and so $S_2 = S = S_1$.

3.3.6 Subsets for Other b , and Symmetry under a $\mathbb{Z}/2$ -action

So far we have established a superset S_1 and a subset S_2 for $b = 1$, and we have noted that $S_1 = S_2$, so $S_2 = S = S_1$. For $b \neq 1$ we have produced a conjectural superset. We also know a very lengthy computation which would verify or refute the conjecture by producing the actual superset. We have not produced a subset S_2 for any $b \neq 1$. The process of producing such an S_2 for SL_3 was a very lengthy and involved proof, and was done in Section 3.1.6. It seems likely that a similar process could be carried out for Sp_4 . We therefore make the following conjecture.

Conjecture 3.3.2. The set $\cup_{(t,e)} S_{(t,e)}$ as (t, e) ranges over all type-edge pairs is in fact equal to S . In other words, $S_1 = S$.




Figure 3.94: Results for $xC_M = D_1$ and $w = r$ or for $xC_M = D_1$ and $w = r^3$




Figure 3.95: Results for $xC_M = D_2$ and $w = r$ or for $xC_M = D_2$ and $w = r^3$




Figure 3.96: Results for $xC_M = D_3$ and $w = r$ or for $xC_M = D_3$ and $w = r^3$




Figure 3.97: Results for $xC_M = D_1$ and $w = r^2$




Figure 3.98: Results for $xC_M = D_2$ and $w = r^2$




Figure 3.99: Results for $xC_M = D_3$ and $w = r^2$

If this conjecture and Conjecture 3.3.1 are both true, the Figures 3.84 through 3.87 represent actual solution sets for their respective σ -conjugacy classes.

In Section 3.1.7, we gave an *a priori* proof of $\mathbb{Z}/3$ -rotational symmetry of the SL_3 solution set S . This proof was based on the existence of a matrix $q \in GL_3(F)$ that acted on A_M by rotating it by 120° around the center of C_M . We have a matrix $q \in GSp_4(F)$ that acts on the main apartment of the building of Sp_4 by flipping it across the (vertical) line of symmetry of C_M . One can prove a proposition exactly analogous to Proposition 3.1.8 in Section 3.1.7 to show that the solution set S for Sp_4 will have symmetry about the vertical line through the middle of C_M . As the rotational symmetry result for SL_3 generalizes easily to a $\mathbb{Z}/n$ -symmetry result for SL_n , so this symmetry result for Sp_4 generalizes to a $\mathbb{Z}/n$ -symmetry result for Sp_{2n} .

3.4 Comments on GSp_4 , PSp_4 , and G_2

If G is GSp_4 or PSp_4 , we have $inv : G(L) \times G(L) \rightarrow \tilde{W} \simeq W_a \ltimes M$, where $M = \mathbb{Z}$ for $G = GSp_4$ and $M = \mathbb{Z}/2$ for $G = PSp_4$. We have $inv(x, y) = (\rho(x^{-1}yC_M), \frac{1}{2}v(\det(x^{-1}y)))$. One can check that if $G = GSp_4$ and $x \in G(L)$ then $v(\det(x)) \in 2\mathbb{Z}$. If $G = PSp_4$ and $x \in G(L)$ then $v(\det(x)) \in 2\mathbb{Z}/4\mathbb{Z} \simeq \mathbb{Z}/2\mathbb{Z}$. The representatives b listed at the beginning of Section 3.3 still represent distinct σ -conjugacy classes in $G(L)$, but there are additional σ -conjugacy classes we will not consider. So if b is one of the matrices at the beginning of Section 3.3, this time considered as an element of $G(L)$, then we ask for a description of the set $\{(\rho(x^{-1}b\sigma(x)C_M), \frac{1}{2}v(\det(x^{-1}b\sigma(x)))) : x \in G(L)\}$. Using methods very similar to those of Section 3.2, we can prove that $\{\rho(x^{-1}b\sigma(x)C_M) : x \in G(L)\} = \{\rho(x^{-1}b\sigma(x)C_M) : x \in Sp_4(L)\}$. It is already clear that $v(\det(x^{-1}b\sigma(x))) = v(\det(b))$ is fixed for fixed b (and equal to 0 for the b we have chosen).

We would also like to make some comments about G_2 . We first note that $W = D_6$, the group of symmetries of a hexagon, and that $W_a = \mathbb{Z}^2 \ltimes W$. We ask for a description of $\{inv(x, \sigma(x)) : x \in G_2(L)\}$. Let ϵ and δ be the two standard generators of $\mathbb{Z}^2 \subseteq \mathbb{Z}^2 \ltimes W \simeq W_a$, where ϵ and δ act on A_M by translating it so that C_M goes, respectively, to the chambers marked with an ϵ and a δ in Figure 3.100.

Let $r \in W$ be the element that rotates A_M counterclockwise by 60° around v_M , and let $f \in W$ be the element that flips A_M about the horizontal line through v_M . We know that ϵ, δ, r and f generate W_a , ϵ and δ commute, and $rf = fr^5$. One can check that $r\epsilon = \epsilon\delta r$, $r\delta = \epsilon^{-1}r$, $f\epsilon = \epsilon f$, and $f\delta = \epsilon^{-1}\delta^{-1}f$.

Using a similar process to that used in Sections 3.1.4 and 3.3.4, if $a = \epsilon^s \delta^t$ and $w \in W$, and if $(aw)^l = 1$ in W_a for some l , then $aw \in \{inv(x, \sigma(x)) : x \in G_2(L)\}$. We get:

Lemma 3.4.1. For any integers s, t and any $1 \leq n \leq 5$, $\epsilon^s \delta^t r^n \in \{inv(x, \sigma(x)) : x \in G_2(L)\}$.




Figure 3.100: The translations ϵ and δ of A_M for G_2

Proof. Just simplify $(\epsilon^s \delta^t r^n)^6$ for each $n = 1, 2, 3, 4, 5$ using the rules listed in the previous paragraph. For instance, if $n = 1$, then $(\epsilon^s \delta^t r)^6 = (\epsilon^s \delta^t r)(\epsilon^s \delta^t r)(\epsilon^s \delta^t r)^4 = (\epsilon^s \delta^t)(\epsilon^{s-t} \delta^s r^2)(\epsilon^s \delta^t r)^4 = (\epsilon^s \delta^t)(\epsilon^{s-t} \delta^s)(\epsilon^{-t} \delta^{s-t})(\epsilon^{-s} \delta^{-t})(\epsilon^{-s+t} \delta^{-s})(\epsilon^t \delta^{-s+t}) r^6 = 1$. $\square$

One could also create lemmas for G_2 analogous to Lemma 3.1.4 and Lemma 3.1.5 in a similar way.

These results give a starting point for the subset S_2 of the solution set S for G_2 with $b = 1$. One could enlarge S_2 using methods similar to those in Section 3.1.5 and Section 3.3.5. One could produce a superset $S_1 \supseteq S$ by folding galleries in an analogous way to Section 3.1.2 and Section 3.3.2. This has not been done, so we cannot say for sure if the resulting S_1 and S_2 are equal, although we guess that they would be.

We also note that G_2 has no analogue to the rotational invariance developed for SL_3 in Section 3.1.7, or the flip invariance developed for Sp_4 in Section 3.3.6, although the general results that will be developed in Section 3.5 will hold for G_2 .

3.5 Invariance Properties of Solution Sets

In this section we prove some results that give invariance properties of the solution set S . These results hold true if S is the solution set $\{inv(x, b\sigma(x)) : x \in G(L)\}$ for any simply-connected group G and any σ -conjugacy class b . We apply the invariance results of this section to SL_3 , $b = 1$ under the assumption that one knows whether $X_w(1\sigma)$ is empty or non-empty only for any w in a certain subset of $W_a = \tilde{W}$. We will see that knowledge on whether $X_w(b\sigma)$ is empty can be obtained on a much larger class of w .

Let G be a simple simply-connected group, let W_a be its affine Weyl group, and let A_M be the main apartment of its building. Suppose G has rank n . Let C_M be the main chamber and let v_M be the main vertex. Let $L_1, \dots, L_n$ be the hyperplanes in A_M that contain v_M and that intersect C_M in an $n - 1$ dimensional simplex. Let L_{n+1} be the hyperplane in A_M that intersects C_M in an $n - 1$ dimensional simplex, but that does not contain v_M . Let s_i be reflection of A_M about L_i . Then the s_i generate W_a as a Coxeter group.

If D is a chamber in A_M , define $p_i D$ to be the chamber obtained by reflecting D about the wall $\tilde{L}_i$, where $\tilde{L}_i$ is parallel to L_i , and intersects D in an $n - 1$ dimensional simplex. So p_i is a map from the set of chambers in A_M to itself. The elements $s_i \in W_a$ are also maps of this kind, and as such we can state the following lemma:

Lemma 3.5.1. If $i \neq j$ then $p_i s_j = s_j p_k$ for some k . We also have $p_i s_i = s_i p_i$.

This can be easily checked, and is valuable because a gallery in A_M starting at C_M is just a sequence of the p_i . We can now prove:

Lemma 3.5.2. If $w \in W_a$ then there is a one-to-one correspondence between Coxeter expansions of w (using the s_i) and galleries from C_M to wC_M . A Coxeter expansion of length m corresponds to a length $m + 1$ gallery, so minimality is preserved under the correspondence.

Proof. If $w = s_{i_1} \cdots s_{i_m}$ then $wC_M = s_{i_1} \cdots s_{i_m} C_M = s_{i_1} \cdots s_{i_{m-1}} p_{i_m} C_M = p_{k_m} s_{i_1} \cdots s_{i_{m-1}} C_M$, where the last equality is achieved by using Lemma 3.5.1 $m - 1$ times. We then proceed: $p_{k_m} s_{i_1} \cdots s_{i_{m-1}} C_M = p_{k_m} s_{i_1} \cdots s_{i_{m-2}} p_{i_{m-1}} C_M = p_{k_m} p_{k_{m-1}} s_{i_1} \cdots s_{i_{m-2}} C_M = \cdots = p_{k_m} p_{k_{m-1}} \cdots p_{k_1} C_M$. Note that length is preserved, and the process is reversible. $\square$

Let $l(w)$ denote the length of $w \in W_a$ as an element of the Coxeter group W_a . Then we have the following results. Note that Proposition 3.5.3 is the affine analogue of some parts of the proof of Theorem 1.6 in [3].

Proposition 3.5.3. If $s = s_i$ for some i and if $l(sws) = l(w)$ then there exists a bijective map from $X_w(b\sigma)$ to $X_{sws}(b\sigma)$.

Proof. We define P to be the parallelogram spanned by C_M and wC_M (so it is the intersection of all apartments containing C_M and wC_M , or, alternatively, the union of all minimal galleries from C_M to wC_M). Because of the fact that $l(sws) = l(w)$, one can show that exactly one of wsC_M and sC_M is in P . If $x \in X_w(b\sigma)$, then let $g \in I$ be such that $gx^{-1}b\sigma(x)C_M = wC_M$. Consider $xg^{-1}P$, which contains $xg^{-1}C_M = xC_M$ and $xg^{-1}wC_M = b\sigma(x)C_M$.

Case 1: $sC_M \subseteq P$. Then let $y = xg^{-1}s$, so $yC_M = xg^{-1}sC_M$. Since yC_M is adjacent to xC_M , $b\sigma(y)C_M$ is adjacent to $b\sigma(x)C_M$, and not a part of $xg^{-1}P$. One can now see that $\text{inv}(y, b\sigma(y)) = swsC_M$.

Case 2: $wsC_M \subseteq P$. Then let $yC_M = \sigma^{-1}b^{-1}xg^{-1}wsC_M$, so we have that $b\sigma(y)C_M = xg^{-1}wsC_M$. Since wsC_M is adjacent to wC_M , $xg^{-1}wsC_M$ is adjacent to $b\sigma(x)C_M$, so yC_M is adjacent to xC_M along the edge $xg^{-1}e$, where e is the edge of adjacency of C_M and sC_M . One can now see that $\text{inv}(y, b\sigma(y)) = swsC_M$.

Now let $\Gamma_1 : X_w(b\sigma) \rightarrow X_{sws}(b\sigma)$ be defined so that $\Gamma_1(xC_M) = yC_M$. I claim that this is a bijection. To check this, note that the roles of w and sws in the construction of Γ_1 were symmetric, so by replacing each with the other we get a map $\Gamma_2 : X_{sws}(b\sigma) \rightarrow X_w(b\sigma)$. One can check that Γ_1 and Γ_2 are inverses. $\square$

Proposition 3.5.4. If $s = s_i$ for some i and if $l(w) > l(sws)$ (and so $l(sws) = l(w) - 2$) then there is a surjective map from $X_w(b\sigma)$ to $X_{sws}(b\sigma) \cup X_{sw}(b\sigma)$.

Proof. Again, let P be the parallelogram spanned by C_M and wC_M . The fact that $l(w) > l(sws)$ means that both wsC_M and sC_M are in P . As in the previous proposition, if $x \in X_w(b\sigma)$, let $g \in I$ be such that $gx^{-1}b\sigma(x)C_M = wC_M$. Let e be the edge of adjacency between wC_M and wsC_M , and consider $xg^{-1}P$. One can see that $b\sigma(xg^{-1}s)C_M$ is adjacent to $b\sigma(x)C_M = xg^{-1}wC_M$ via $xg^{-1}e$, but we may have $b\sigma(xg^{-1}s)C_M = xg^{-1}wsC_M$ or we may have $b\sigma(xg^{-1}s)C_M \neq xg^{-1}wsC_M$ (at least *a priori*, either the equality or the inequality could possibly hold, and it will turn out that both actually do arise).


Case 1: $b\sigma(xg^{-1}s)C_M = xg^{-1}wsC_M$. Then if we let $y = xg^{-1}s$, we have that $\text{inv}(y, b\sigma(y)) = \text{inv}(xg^{-1}s, xg^{-1}ws) = \text{inv}(s, ws) = \text{inv}(1, sws) = sws$, so $y \in X_{sws}(b\sigma)$.

Case 2: $b\sigma(xg^{-1}s)C_M \neq xg^{-1}wsC_M$. Then if $y = xg^{-1}s$, we have $\text{inv}(y, b\sigma(y)) = \text{inv}(s, w) = \text{inv}(1, sw) = sw$, so $y \in X_{sw}(b\sigma)$.

So we have a map $\Gamma : X_w(b\sigma) \rightarrow X_{sws}(b\sigma) \cup X_{sw}(b\sigma)$ where $\Gamma(xC_M) = yC_M$. We must show that Γ is surjective. Take $z \in X_{sw}(b\sigma)$, so $\text{inv}(z, b\sigma(z)) = sw$. So $\rho_{sC_M}(sz^{-1}b\sigma(z)C_M) = wC_M$, where ρ_{sC_M} is the retraction of $\mathcal{B}_\infty$ onto A_M centered at sC_M . So there exists $g \in sIs$ such that $gsz^{-1}b\sigma(z)C_M = wC_M$. Consider $zsg^{-1}P$, which contains zC_M and $b\sigma(z)C_M = zsg^{-1}wC_M$. Note that if e is the edge of adjacency of C_M and sC_M , then zC_M contains $zsg^{-1}e$. Choose a chamber xC_M containing e , but not equal to zC_M . Then $b\sigma(x)C_M$ is adjacent to $b\sigma(z)C_M$. We may assume without loss of generality that $b\sigma(x)C_M \not\subseteq zsg^{-1}P$ (we can arrange this by choosing an appropriate xC_M). Then $x \in X_w(b\sigma)$ and $\Gamma(xC_M) = zC_M$.

If $z \in X_{sws}(b\sigma)$, then $\rho_{sC_M}(sz^{-1}b\sigma(z)C_M) = wsC_M$. So there exists $g \in sIs$ such that $gsz^{-1}b\sigma(z)C_M = wsC_M$. Consider $zsg^{-1}P$, which contains zC_M and $b\sigma(z)C_M = zsg^{-1}wsC_M$. Let e be the edge between C_M and sC_M . We know $zsg^{-1}e \subseteq zC_M$, and $b\sigma(e) = zsg^{-1}we$. So choose xC_M containing e and not in P . Then $b\sigma(x)C_M$ contains $b\sigma(e)$. We can require without loss of generality that $b\sigma(x)C_M \not\subseteq P$. Then $\Gamma(xC_M) = zC_M$. This proves surjectivity. $\square$

Corollary 3.5.5. If $l(sws) > l(w)$ (so $l(sws) = l(w) + 2$) then there is a surjective map from $X_{sws}(b\sigma)$ to $X_w(b\sigma) \cup X_{ws}(b\sigma)$.

Figure 3.101: Region R

Proof. Just apply the previous proposition with sws in place of w . □

Corollary 3.5.6. If $l(sw) > l(w)$, $l(ws) < l(w)$ and $X_w(b\sigma) \neq \emptyset$ then $X_{sw}(b\sigma) \neq \emptyset$. If $l(sw) < l(w)$, $l(ws) > l(w)$, and $X_w(b\sigma) \neq \emptyset$, then $X_{ws}(b\sigma) \neq \emptyset$.

Proof. This follows from the two propositions. □

Note that we used this last result repeatedly in Section 3.1.5 and Section 3.3.5 for SL_3 and Sp_4 .

These results are valuable because they can be used on a wide variety of groups G and σ -conjugacy classes b to increase partial information on the nature of the solution set $\{inv(x, b\sigma(x)) : x \in G(L)\}$. To illustrate the possible utility of the above propositions, we consider the hypothetical case that for SL_3 , $b = 1$, one knows for any w in region R on Figure 3.101 whether $X_w(1\sigma)$ is or is not empty. So we assume that part of Figure 3.27, the solution set for SL_3 , $b = 1$, is given. In this case, one can repeatedly apply the results of this section, together with the rotational invariance result of Section 3.1.7, to the point where it is known whether $X_w(b\sigma)$ is empty or non-empty for each w corresponding to a chamber shaded in Figure 3.102. We note that this represents nearly complete information about the solution set of SL_3 , $b = 1$, since nearly every chamber in that figure is shaded. The results of this section could be applied with the same level of effectiveness if $b \neq 1$, or for Sp_4 . Our results could even be applied to other groups, including higher rank groups.

Also note that Sections 3.1.5 and 3.3.5 were essentially just Corollary 3.5.6 applied to the partial information of Sections 3.1.4 and 3.3.4.




Figure 3.102: Understood region after applying invariance properties to region R

REFERENCES

- [1] Bruhat, F. and Tits, J. *Groupes reductifs sur un corps local*, Inst. Hautes Etudes Sci. Publ. Math. 41 (1972), 5–251.
- [2] Bruhat, F. and Tits, J. *Groupes reductifs sur un corps local II*, Inst. Hautes Etudes Sci. Publ. Math. 60 (1984), 197–376.
- [3] Deligne, P. and Lusztig, G. *Representations of Reductive Groups Over Finite Fields*, Annals of Mathematics 103 (1976), 103–161.
- [4] Garrett, P. *Buildings and Classical Groups*, Chapman and Hall (1997).
- [5] Kneser, M. *Chapter 10 of Algebraic Number Theory*, edited by Cassels, J.W.S., and Frohlich, A., Academic Press, (1967).
- [6] Kottwitz, R. *Isocrystals with Additional Structure*, Compositio Mathematica 56 (1985), 201–220.
- [7] Kottwitz, R. *Isocrystals with Additional Structure II*, Compositio Mathematica 109 (1997), 255–339.
- [8] Kottwitz, R. *Orbital Integrals on GL_3* , American Journal of Mathematics 102 (1980), 327–384.
- [9] Kottwitz, R. and Rapoport, M. *On the Existence of F -crystals*, (2002).
- [10] Manin, Y. *The Theory of Commutative Formal Groups Over Fields of Finite Characteristic*, Russian Mathematical Survey 18 #6 (1963), 1–81.
- [11] Rapoport, M. *A Positivity Property of the Satake Isomorphism*, preprint.
- [12] Serre, J.-P. *Galois Cohomology*, Springer Verlag (1997).
- [13] Serre, J.-P. *Local Fields*, Springer Verlag (1979).